1899

Mafic Inclusions, Aggregates, and Dikes in Granitoid Rocks, Central Sierra Nevada Batholith, California—Analytic Data

U.S. GEOLOGICAL SURVEY BULLETIN 1899

READEX FICHE best copy available

I 19.3:

AVAILABILITY OF BOOKS AND MAPS OF THE U.S. GEOLOGICAL SURVEY

Instructions on ordering publications of the U.S. Geological Survey, along with prices of the last offerings, are given in the current-year issues of the monthly catalog "New Publications of the U.S. Geological Survey." Prices of available U.S. Geological Survey publications released prior to the current year are listed in the most recent annual "Price and Availability List." Publications that are listed in various U.S. Geological Survey catalogs (**see back inside cover**) but not listed in the most recent annual "Price and Availability List" are no longer available.

Prices of reports released to the open files are given in the listing "U.S. Geological Survey Open-File Reports," updated monthly, which is for sale in microfiche from the U.S. Geological Survey, Books and Open-File Reports Section, Federal Center, Box 25425, Denver, CO 80225. Reports released through the NTIS may be obtained by writing to the National Technical Information Service, U.S. Department of Commerce, Springfield, VA 22161; please include NTIS report number with inquiry.

Order U.S. Geological Survey publications **by mail** or **over the counter** from the offices given below.

BY MAIL

Books

Professional Papers, Bulletins, Water-Supply Papers, Techniques of Water-Resources Investigations, Circulars, publications of general interest (such as leaflets, pamphlets, booklets), single copies of Earthquakes & Volcanoes, Preliminary Determination of Epicenters, and some miscellaneous reports, including some of the foregoing series that have gone out of print at the Superintendent of Documents, are obtainable by mail from

U.S. Geological Survey, Books and Open-File Reports
Federal Center, Box 25425
Denver, CO 80225

Subscriptions to periodicals (Earthquakes & Volcanoes and Preliminary Determination of Epicenters) can be obtained ONLY from the

Superintendent of Documents
Government Printing Office
Wa.hington, D.C. 20402

(Check or money order must be payable to Superintendent of Documents.)

Maps

For maps, address mail orders to

U.S. Geological Survey, Map Distribution
Federal Center, Box 25286
Denver, CO 80225

Residents of Alaska may order maps from

Alaska Distribution Section, U.S. Geological Survey,
New Federal Building - Box 12
101 Twelfth Ave., Fairbanks, AK 99701

OVER THE COUNTER

Books

Books of the U.S. Geological Survey are available over the counter at the following Geological Survey Public Inquiries Offices, all of which are authorized agents of the Superintendent of Documents:

- **WASHINGTON, D.C.**--Main Interior Bldg., 2600 corridor, 18th and C Sts., NW.
- **DENVER, Colorado**--Federal Bldg., Rm. 169, 1961 Stout St.
- **LOS ANGELES, California**--Federal Bldg., Rm. 7638, 300 N. Los Angeles St.
- **MENLO PARK, California**--Bldg. 3 (Stop 533), Rm. 3128, 345 Middlefield Rd.
- **RESTON, Virginia**--503 National Center, Rm. 1C402, 12201 Sunrise Valley Dr.
- **SALT LAKE CITY, Utah**--Federal Bldg., Rm. 8105, 125 South State St.
- **SAN FRANCISCO, California**--Customhouse, Rm. 504, 555 Battery St.
- **SPOKANE, Washington**--U.S. Courthouse, Rm. 678, West 920 Riverside Ave..
- **ANCHORAGE, Alaska**--Rm. 101, 4230 University Dr.
- **ANCHORAGE, Alaska**--Federal Bldg. Rm. E-146, 701 C St.

Maps

Maps may be purchased over the counter at the U.S. Geological Survey offices where books are sold (all addresses in above list) and at the following Geological Survey offices:

- **ROLLA, Missouri**--1400 Independence Rd.
- **DENVER, Colorado**--Map Distribution, Bldg. 810, Federal Center
- **FAIRBANKS, Alaska**--New Federal Bldg., 101 Twelfth Ave.

Mafic Inclusions, Aggregates, and Dikes in Granitoid Rocks, Central Sierra Nevada Batholith, California—Analytic Data

By BERNARD BARBARIN, F.C.W. DODGE, R.W. KISTLER, and P.C. BATEMAN

U.S. GEOLOGICAL SURVEY BULLETIN 1899

DEPARTMENT OF THE INTERIOR
MANUEL LUJAN, JR., Secretary

U.S. GEOLOGICAL SURVEY
Dallas L. Peck, Director

Any use of trade, product, or firm names in this publication is for descriptive purposes only and does not imply endorsement by the U.S. Government

UNITED STATES GOVERNMENT PRINTING OFFICE, WASHINGTON : 1989

For sale by the Books and
Open-File Reports Section,
U.S. Geological Survey
Federal Center, Box 25425
Denver, CO 80225

Library of Congress Cataloging-in-Publication Data

Mafic inclusions, aggregates, and dikes in granitoid rocks, Central Sierra Nevada batholith, California—analytic data / by Bernard Barbarin . . . [et al.]
p. cm. — (U.S. Geological Survey bulletin ; 1899)
Includes bibliographical references.
Supt. of Docs. no.: I 19.3; 1899
1. Granite—California—Sierra Nevada Region—Inclusions. 2. Rocks, Ultrabasic—California—Sierra Nevada Region. 3. Geology—California—Sierra Nevada Region. I. Barbarin, Bernard. II. Series.
QE75.B9 no. 1899 557.3 s—dc20 89-600250
[QE462.G7] [552'.3] CIP

CONTENTS

Abstract 1
Introduction 1
Occurrence and distribution 2
Sampling and analytical methods 2
Host rocks 3
Aggregates 3
Inclusions and dike rocks 3
Mineral compositions 3
References cited 3

FIGURES

1. Simplified geologic map of the Sierra Nevada between latitudes 37° and 38° N. 7
2. Modal plots of mafic inclusions, aggregates, dikes, and their host granitoids 8
3. Normative plots of mafic inclusions, aggregates, dikes, and their host granitoids 8
4. Plot of mutually exclusive normative constituents of mafic inclusions, aggregates, dikes, and their host granitoids 9

TABLES

1. Location and rock unit of collected samples 10
2. Modal analyses, in volume percent, of mafic inclusions, aggregates, dikes, and their host granitoids and associated gabbroid 11
3. Chemical and trace-element analyses of mafic inclusions, aggregates, dikes, and their host granitoids, associated gabbroid, and aplite 13
4. CIPW norms of mafic inclusions, aggregates, dikes, and their host granitoids, associated gabbroid, and aplite 21
5. Average chemical analyses, in weight percent, of biotite in mafic inclusions, dikes, aggregates, and their host granitoids 24
6. Average chemical analyses, in weight percent, of hornblende in mafic inclusions, dikes, aggregates, and their host granitoids and associated gabbroid 26
7. Rb-Sr isotopic data on mafic inclusions, dikes, aggregates, and their host granitoids 28
8. Sm-Nd isotopic data on mafic inclusions, aggregates, and associated rocks 29

Mafic Inclusions, Aggregates, and Dikes in Granitoid Rocks, Central Sierra Nevada Batholith, California—Analytic Data

By Bernard Barbarin,[1] F.C.W. Dodge, R.W. Kistler, *and* P.C. Bateman

ABSTRACT

Analytic data are reported herein for a comprehensively studied suite of samples of mafic inclusions, aggregates or schliere, and dikes and their host granitoids from 12 localities in the central Sierra Nevada. Modes are given for 61 of the samples, and major-element oxides and selected trace-element concentrations are tabulated for 81. Chemical analyses are presented for coexisting hornblendes and biotites in the inclusions, dikes, aggregates, and their host granitoids.

INTRODUCTION

During the past 25 years, analytic data on granitoid rocks of the central Sierra Nevada have been included in numerous reports of the U.S. Geological Survey by geologists studying 15-minute quadrangles in the region between latitudes 37° and 38° N. (for example, Bateman, 1965a; Bateman and Busacca, 1983; Bateman and others, 1984; Bateman and Lockwood, 1970, 1976; Bateman and Wones, 1972a; Dodge and Calk, 1986). These data are typical of large parts of mapped granitoid bodies of the Sierra Nevada batholith. However, many of these large granitoid units contain mafic inclusions, streaky aggregates, and dikes that generally have not been collected and analyzed. This report provides modal, chemical, and isotopic data on such rocks and their hosts, and it provides chemical data on constituent hornblendes and biotites from a dozen localities selected from four mapped granitoid units. Limited additional data were obtained on samples taken from other localities; however, these data are incomplete and are not included herein.

Recent reports on the Sierra Nevada have referred to dark, fine-grained clots as "mafic inclusions" or "mafic enclaves" to avoid any genetic implications. Similar inclusions have been called basic concretions, autoliths, xenoliths, magmatic enclaves, or other names that usually have a genetic connotation. These inclusions, although variable in size, have well-defined shapes, are generally in sharp contact with their hosts, and typically are distinctly finer grained than their hosts. Less-compact aggregates of mafic minerals commonly have diffuse, irregular outlines and may show little difference in grain size compared with their hosts. Both gradations and abrupt discontinuities are present between aggregates and host granitoids, but only abrupt discontinuities are present between inclusions and host granitoids. Streaky aggregates consisting principally of mafic minerals are locally associated with the inclusions. Although all elongate mafic aggregates or segregations, including elongate inclusions, could be included under the term "schlieren," geologists in the Sierra Nevada have followed the usage of Cloos (1936), who restricted the term to composite layers within plutons that grade from mafic, fine-grained rock to more felsic, coarser grained rock. Therefore, we have simply called all irregular, streaky concentrations of mafic minerals, whether or not layered, aggregates. Mafic dikes occur as both continuous and fragmental sheetlike or tabular bodies that commonly cut primary foliations, whereas inclusions commonly define the foliations.

A Ph.D. thesis study by Adolf Pabst over 60 years ago resulted in the publication of his often-cited classic paper on inclusions in Sierra Nevada granitoids (Pabst, 1928). More recently, these mafic inclusions have been the prime subject of a few studies (Link, 1968; Reid and others, 1983; Furman and Spera, 1985), but they have also been mentioned in several broader investigations (Bateman, 1965b, in press; Loomis, 1983; Moore, 1963; Noyes and others, 1983; Piwinskii, 1968; Sherlock and Hamilton, 1958). The present study began when P.C. Bateman and F.C.W. Dodge started collecting inclusions and associated rocks as an adjunct to regional studies in the central Sierra Nevada. Subsequently, Bernard Barbarin collected additional samples from localities in regions mapped by Bateman, Dodge, and other U.S. Geological Survey geologists (Bateman, 1965a; Bateman and Busacca, 1982; Bateman and others, 1971; Bateman and Wones, 1972b; Dodge and Calk, 1987; Lockwood

Manuscript approved for publication June 14, 1989.

[1]Université de Paris-Sud, Laboratoire de Pétrographie–Volcanologie, Bât. 504, F–91405 Orsay Cedex, France.

and Bateman, 1976) and conducted an intensive study of all collected samples. Barbarin's work was done while he was the recipient of a "Lavoisier" grant from the French government and with U.S. Geological Survey support.

OCCURRENCE AND DISTRIBUTION

Mafic inclusions, although widespread, are almost exclusively restricted to hornblende-bearing granitoids and are most common in felsic tonalites and mafic granodiorites. Commonly they are uniformly distributed over broad areas, although local concentrations are not uncommon. Local accumulations mainly occur near contacts of one granitoid body with another, but they are also found far from contacts. In compositionally zoned granitoid bodies, the abundance of inclusions commonly varies sympathetically with the mafic mineral content of the host, generally decreasing from margins to interiors of granitoid bodies. Moore (1963, p. 119) showed that thicker plutons have smaller proportions of inclusions than thinner ones, regardless of mafic mineral content. Inclusion abundance and age are not apparently correlative; inclusions are present in rocks ranging from Late Triassic to Late Cretaceous, the entire age span of granitoids in the central Sierra Nevada.

Unlike mafic inclusions, streaky aggregates of mafic minerals are locally concentrated in granitoids of all compositions; the occurrence of these aggregates rarely exceeds such local concentrations. Zones of aggregates are most common near contacts between granitoids of markedly different compositions or near contacts of granitoids with country rocks, but they also occur within plutons, far from external contacts. Mafic inclusions are typically associated with the aggregates, particularly near contacts and where the host granitoid composition is appropriate.

Mafic dikes occur sporadically, singly, and in small groups as well as in large, pervasive multitudes, as in the Independence and Sonora dike swarms (Moore and Hopson, 1961; Schweickert and Bogen, 1983). Dikes may be common in one part of a pluton but sparse or nonexistent elsewhere, and they are totally absent in many plutons, particularly those that are felsic. Typically, individual dikes are a few centimeters to a few meters thick and generally are vertical or dip gently, as along Stevenson Creek, a tributary of the San Joaquin River. Most dikes have sharp walls that cut across the foliation of enclosing rocks, although some appear to have been deformed and cleaved along with their hosts. Rarely, as at Hetch Hetchy, the dikes are disrupted or broken up into inclusion-like bodies, although such dikes maintained their original continuity. Some dikes are net veined with felsic aplitic material that may compose a large portion of the dike. Contacts between the felsic and mafic parts of these dikes are typically crenulate.

SAMPLING AND ANALYTICAL METHODS

Various numbering schemes were used when samples were originally collected. For consistency, samples were redesignated using a uniform system. The first two letters reflect the sample locality (see Table 1). These capital letters are followed by the number of samples collected from the locality; a lowercase letter following a number indicates whether more than one sample, and how many, have been collected from a single dike or inclusion. The alphanumeric string is followed by a dash and a capital letter designating sample type: H, host granitoid; A, aggregate; D, dike; I, inclusion; G, gabbroid; AP, aplite. For some samples this letter is followed by "m" or "c" indicating the margin or core of an inclusion or by "x" or "i" indicating exterior or interior of a dike. Thus, MHb9b-Ic refers to the core of an inclusion collected from Camp 61.

Location and the distribution of rock units from which samples were taken are given in table 1 and shown in figure 1. The size of sampled outcrop varies but typically is within an area of tens of square meters or less.

Modal analyses of medium- to coarse-grained rocks were determined by combining the point counts of selectively stained rock slabs (Norman, 1974) with those of thin sections; modal analyses of fine-grained rocks were determined from thin sections only. Sixty-one modal analyses, including 33 of inclusions, 13 of host rocks, 7 each of mafic aggregates and dike rocks, and 1 gabbro, are given in table 2. Chemical and trace-element analyses of 39 mafic inclusions, 15 host rocks, 14 dike rocks, 10 mafic aggregates, 1 aplite, 1 gabbro, and 1 hornfels inclusion are given in table 3; CIPW norms computed from the chemical analyses are given in table 4. All samples were thoroughly examined and carefully prepared to ensure that only the designated rock type was contained in the analyzed material.

Electron microprobe analyses of biotite and hornblende, 67 of each mineral, are given in tables 5 and 6. Each quoted analysis represents the mean of three to nine individual analyses of mineral grains from a specific rock type within a sample. Whenever possible, microprobe mounts were prepared to include parts of host rock, inclusion, and aggregate so that minerals from all the rock types could be analyzed from a single mount. Separate analyses are presented for hornblendes and biotites from cores or darker areas of a few inclusions.

The modal amounts of plagioclase, mafic plus accessory minerals, and quartz plus potassium feldspar, and the amounts of plagioclase, quartz, and potassium feldspar recalculated to 100 percent are plotted on ternary diagrams (fig. 2). Similarly, percentage of normative albite plus anorthite, quartz plus orthoclase, and all other constituents, and the percentage of normative albite plus anorthite, quartz, and orthoclase recalculated to 100 percent are plotted (fig. 3). In addition, the

amounts of the mutually exclusive normative constituents olivine or quartz and diopside or corundum, indicators of silica and alumina under- or oversaturation, are shown in figure 4.

A total of 65 strontium and 11 neodymium isotopic analyses of host rock, inclusions, mafic aggregates, and dike rocks are given in tables 7 and 8, respectively. Errors reported are within two standard deviations of the mean. Values of $^{87}Sr/^{86}Sr$ are normalized to a value of $^{86}Sr/^{88}Sr=0.1194$ and replicate analyses of $SrCO_3$ standard NBS 987 give a value of $^{87}Sr/^{86}Sr=0.710236\pm3$. Values of $^{143}Nd/^{144}Nd$ are normalized to a value of $^{146}Nd/^{144}Nd=0.7219$ and replicate analyses of USGS rock standard BCR–1 give a value of $^{143}Nd/^{144}Nd=0.512642$.

HOST ROCKS

Compositions of host rocks are restricted to but span the entire granodiorite and tonalite fields. The amount of mafic minerals generally ranges between 10 and 25 percent, although it is greater for a few of the samples. Silica contents range between 60 and 70 weight percent and potassium-oxide contents range between 1.7 and 4.0 percent. Norms of the host rocks are more closely clustered on the ternary diagrams than are the modes. All the granitoid hosts are quartz normative and most are weakly alumina saturated, falling within the field defined by norms of other Sierra Nevada plutonic rocks. Trace-element contents are similar to those reported by Dodge and others (1982) for hornblende-bearing Sierra Nevada granitoid rocks.

AGGREGATES

Because of their inherent inhomogeneity, analytical data of the mafic aggregates or schliere are difficult to evaluate. Commonly, the analytical data overlap those of the other groups and all plots show considerable scatter.

INCLUSIONS AND DIKE ROCKS

Modes of inclusions tend to plot along the plagioclase-quartz join of the quartz-potassium feldspar-plagioclase diagram within the diorite, quartz diorite, and tonalite fields. Mafic and accessory minerals make up less than 25 to more than 60 volume percent of the inclusions but generally are between 30 and 55 percent. On the triangular plot of all constituents, the modes of host rocks and inclusions form an arcuate field with mafic inclusions forming the end of the field closer toward mafic plus accessory minerals and their hosts forming the end closer toward quartz plus alkali feldspar. Although the field is continuous, there is little overlap between the two groups. A similar relation is shown by norms, although norms are displaced toward the quartz plus orthoclase corner, which reflects the content of potassium in biotite in computing normative orthoclase. Also, at least in part because of biotite, the inclusions tend to lie on or near the orthoclase-albite plus anorthite join of the quartz-albite plus anorthite-orthoclase diagram, even though all the inclusions contain at least minor or trace amounts of modal quartz. In fact, about half the inclusions are olivine normative. Only two quartz-normative inclusions contain normative corundum. Silica contents range from 46 to 63 weight percent and generally are between 50 and 60 percent, and potassium-oxide contents range from 0.72 to 3.93 percent but commonly are between 1.5 and 2.5 percent. Of the trace elements, the "compatible" or transition elements Sc, V, Cr, and Co are concentrated in inclusions relative to host rocks. Concentration of Zr is about the same in inclusions as in host rocks. Concentration of Ba tends to be lower in inclusions, but concentration of Rb is generally greater. Rare-earth elements are commonly less fractionated in inclusions than in the host granitoids, and La/Yb ratios of the inclusions are about half those of the hosts, although total rare-earth-element contents are greater in the inclusions.

Generally, the analytic data on samples of mafic dikes are similar to those of the mafic inclusions, particularly the more quartz-rich inclusions.

MINERAL COMPOSITIONS

Biotite and hornblende in mafic inclusions, dikes, and aggregates have the same chemical composition as in the contiguous granitoids, regardless of differences in grain size, morphology, crystal habit, or location within the inclusions. Atomic Fe/Fe+Mg ratios clearly show this relation. Where compositions of the mafic minerals change from one locality to another within a pluton, parallel changes occur within the inclusions. These compositional variations also occur in biotite and hornblende from closely spaced inclusion-granitoid pairs from the same locality.

REFERENCES CITED

Bateman, P.C., 1965a, Geologic map of the Blackcap Mountain quadrangle, Fresno County, California: U.S. Geological Survey Geologic Quadrangle Map GQ-428, scale 1:62,500.

———1965b, Geology and tungsten mineralization of the Bishop district, California: U.S. Geological Survey Professional Paper 470, 208 p.

———in press, Constitution and genesis of the central part of the Sierra Nevada batholith, California: U.S. Geological

Survey Professional Paper 1483.

Bateman, P.C., and Busacca, A.J., 1982, Geologic map of the Millerton Lake quadrangle, west-central Sierra Nevada, California: U.S. Geological Survey Geologic Quadrangle Map GQ-1548, scale 1:62,500.

———1983, Millerton Lake quadrangle, west-central Sierra Nevada, California: U.S. Geological Survey Professional Paper 1261, 20 p.

Bateman, P.C., Dodge, F.C.W., and Bruggman, P.E., 1984, Major oxide analyses, CIPW norms, modes, and bulk specific gravities of plutonic rocks from the Mariposa 1° by 2° sheet, central Sierra Nevada, California: U.S. Geological Survey Open-File Report 84–162, 50 p.

Bateman, P.C., and Lockwood, J.P., 1970, Kaiser Peak quadrangle, central Sierra Nevada, California—analytic data: U.S. Geological Survey Professional Paper 644-C, p. C1-C15.

———1976, Shaver Lake quadrangle, central Sierra Nevada, California—analytic data: U.S. Geological Survey Professional Paper 774-D, p. D1-D20.

Bateman, P.C., Lockwood, J.F., and Lydon, P.A., 1971, Geologic map of the Kaiser Peak quadrangle, central Sierra Nevada, California: U.S. Geological Survey Geologic Quadrangle Map GQ-894, scale 1:62,500.

Bateman, P.C., and Wones, D.R., 1972a, Huntington Lake quadrangle, central Sierra Nevada, California—analytic data: U.S. Geological Survey Professional Paper 724-A, p. A1-A18.

———1972b, Geologic map of the Huntington Lake quadrangle, central Sierra Nevada, California: U.S. Geological Survey Geologic Quadrangle Map GQ-987, scale 1:62,500.

Cloos, Ernst, 1936, Der Sierra-Nevada-Pluton in Californien: Neues Jahrbuch für Mineralogie, Geologie und Palaentologie, supp., v. 76, no. 3, sec. B, p. 355–450.

Dodge, F.C.W., and Calk, L.C., 1986, Lake Eleanor quadrangle, central Sierra Nevada, California—analytic data: U.S. Geological Survey Bulletin 1565, 20 p.

———1987, Geologic map of the Lake Eleanor quadrangle, central Sierra Nevada, California: U.S. Geological Survey Geologic Quadrangle Map GQ-1639, scale 1:62,500.

Dodge, F.C.W., Millard, H.T., Jr., and Elsheimer, H.N., 1982, Compositional variations and abundances of selected elements in granitoid rocks and constituent minerals, central Sierra Nevada batholith, California: U.S. Geological Survey Professional Paper 1248, 24 p.

Domenick, M.A., Kistler, R.W., Dodge, F.C.W., and Tatsumoto, M, 1983, Nd and Sr isotopic study of crustal and mantle inclusions from the Sierra Nevada and implications for batholith petrogenesis: Geological Society of America Bulletin, v. 94, p. 713–719.

Furman, Tanya, and Spera, F.J., 1985, Co-mingling of acid and basic magma with implications for the origin of mafic I-type xenoliths: Field and petrochemical relations of an unusual dike complex at Eagle Lake, Sequoia National Park, California, U.S.A.: Journal of Volcanology and Geothermal Research, v. 24, p. 151–178.

Link, A., 1968, Inclusions in the Half Dome Quartz Monzonite, Yosemite National Park, California: Evanston, Ill., Northwestern University, unpublished Ph.D. thesis.

Lockwood, J.P., and Bateman, P.C., 1976, Geologic map of the Shaver Lake quadrangle, central Sierra Nevada, California: U.S. Geological Survey Geologic Quadrangle Map GQ-1271, scale 1:62,500.

Loomis, A.A., 1983, Geology of the Fallen Leaf Lake 15′ quadrangle, El Dorado County, California: California Division of Mines and Geology Map Sheet 32, scale 1:62,500.

Moore, J.G., 1963, Geology of the Mount Pinchot quadrangle, southern Sierra Nevada, California: U.S. Geological Survey Bulletin 1130, 152 p.

Moore, J.G., and Hopson, C.A., 1961, The Independence dike swarm in eastern California: American Journal of Science, v. 259, p. 241–259.

Norman, M.B., 1974, Improved techniques for selective staining of feldspar and other minerals using amaranth: U.S. Geological Survey Journal of Research, v. 2, p. 73–79.

Noyes, H.J., Frey, F.A., and Wones, D.R., 1983, A tale of two plutons: Geochemical evidence bearing on the origin and differention of the Red Lake and Eagle Peak plutons, central Sierra Nevada, California: Journal of Geology, v. 91, no. 5, p. 487–509.

Pabst, Adolf, 1928, Observations on inclusions in granitic rocks of the Sierra Nevada: University of California Publications, Bulletin of Sciences, v. 17, p. 325–386.

Piwinskii, A.J., 1968, Studies of batholithic feldspars: Sierra Nevada, California: Contributions to Mineralogy and Petrology, v. 17, p. 204–223.

Reid, J.B., Jr., Evans, O.C., and Fates, D.G., 1983, Magma mixing in granitic rocks of the central Sierra Nevada, California: Earth and Planetary Science Letters, v. 66, p. 243–261.

Schweickert, R.A., and Bogen, N.L., 1983, Tectonic transect of Sierran Paleozoic through Jurassic accreted belts: Society of Economic Paleontologists and Mineralogists, Pacific Section, 22 p.

Sherlock, D.G., and Hamilton, W.B., 1958, Geology of the north half of the Mount Abbot quadrangle, Sierra Nevada, California: Geological Society of America Bulletin, v. 69, p. 1245–1267.

Streckeisen, A.L., 1973, Plutonic rocks: Classification and nomenclature recommended by the IUGS subcommission on the systematics of igneous rocks: Geotimes, v. 18, no. 10, p. 26–30.

FIGURES 1–4; TABLES 1–8



Figure 1. Simplified geologic map of the Sierra Nevada between latitudes 37° and 38° N. showing distribution of granitoid rock units and locations from which samples were taken.







Figure 2. Modal plots of mafic inclusions, aggregates, dikes, and their host granitoids. *A*, Plot on plagioclase-quartz-potassium feldspar diagram. *B*, Classification from Streckeisen and others (1973). *C*, Plot on diagram of all modal constituents.





Figure 3. Normative plots of mafic inclusions, aggregates, dikes, and their host granitoids. *A*, Plot on albite + anorthite-quartz-orthoclase diagram. *B*, Plot on diagram of all normative constituents.



Figure 4. Plot of mutually exclusive normative constituents of mafic inclusions, aggregates, dikes, and their host granitoids showing silica and alumina over- or under-saturation. Outlined field includes more than 90 percent of CIPW norms of Sierra Nevada plutonic rocks from Bateman and others (1984).

Table 1. Location and rock unit of collected samples

Map Symbol (fig.1)	Locality	Lat. (N.)	Long. (W.)	Rock unit
HH	Hetch Hetchy	37°56.8'	119°47.2'	Bass Lake Tonalite
NF	North Fork	37°11.6'	119°35.8'	Do.
ML	Meadow Lakes	37°04.4'	119°24.5'	Do.
SC	Stevenson Creek	37°11.2'	119°20.9'	Do.
SQ	Shaver Lake quarry	37°09.3'	119°17.7'	Dinkey Creek Granodiorite
CS	Camp Sierra	37°11.6'	119°15.6'	Do.
MK	Markwood Creek	37°05.8'	119°14.7'	Do.
DR	Deer Creek	37°00.3'	119°04.0'	Do.
WR	Wishon Reservoir	36°59.9'	118°57.6'	Do.
CP	Chinese Peak	37°13.3'	119°09.3'	Granodiorite of Red Lake
MHa	Mono Hot Springs	37°19.5'	119°00.8'	Mount Givens Granodiorite
MHb	Camp 61	37°19.1'	119°03.7'	Do.
CR	Courtright Reservoir	37°04.4'	118°58.0'	Do.

Table 2. Modal analyses, in volume percent, of mafic inclusions, aggregates, dikes, and their host granitoids and associated gabbroid

[Analyses by B. Barbarin. Rock type indicated by letter following hyphen: H, host granitoid; A, aggregate; I, inclusion; D, dike. Letter following rock type: x, exterior; i, interior; m, margin; c, core]

	Bass Lake Tonalite								
	Hetch Hetchy						North Fork		
	Tonalite	Aggregate	Inclusion	Dike rock			Tonalite	Inclusion	
	HH1-H	HH3-A	HH4-I	HH6a-D	HH6b-D	HH6d-D	NF1-H	NF2-I	NF3-I
Quartz	26.6	20.8	2.7	3.9	9.6	20.2	25.9	4.2	1.7
K-feldspar	19.2	0.6	0	0.8	0.1	0.5	9.5	1.4	0
Plagioclase	37.3	50.4	49.9	49.1	47.5	49.4	44.8	50.3	51.0
Biotite	8.7	15.0	21.5	13.3	12.6	16.0	12.3	13.2	18.6
Hornblende	6.3	11.5	23.3	30.0	27.8	12.4	6.7	30.6	28.6
Sphene			1.1	2.1	1.3				
Opaque	1.9	1.7	1.1	0.3	0.3	1.5	0.9	0.3	0.1
Other			0.4	0.5	0.8				
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.1	100.0	100.0

	Bass Lake Tonalite—Continued					Dinkey Creek Granodiorite			
	North Fork—Continued			Meadow Lakes		Shaver Lake quarry			
	Dike rock		Gabbro	Tonalite	Inclusion	Granodiorite		Inclusion	
	NF4a-D	NF4b-D	NF5-G	ML1-H	ML2-I	SQ1-H	SQ2-H	SQ3-I	SQ5-I
Quartz	1.4	0.8	16.8	20.3	3.1	27.1	25.4	4.2	5.8
K-feldspar	0	0	0.4	2.8	0.4	13.9	12.3	0	0.6
Plagioclase	52.3	43.7	43.7	47.8	50.1	40.2	41.3	46.0	47.3
Biotite	9.8	16.9	15.4	15.9	12.1	12.1	12.0	18.0	18.2
Hornblende	32.9	36.2	22.7	11.6	29.6	4.4	7.5	30.6	27.2
Sphene	0.4	0			1.2				
Opaque	2.5	1.7	1.0	1.6	2.5	2.3	1.5	1.2	0.9
Other	0.7	0.7			1.0				
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

	Dinkey Creek Granodiorite—Continued								
	Shaver Lake quarry—Continued						Camp Sierra		
	Inclusion—Continued				Dike rock		Granodiorite	Inclusion	
	SQ6-I	SQ9-I	SQ10-I	SQ11-I	SQ12-Dx	SQ12-Di	CS1-H	CS2-I	CS3-I
Quartz	1.2	17.2	5.2	0.8	13.4	11.9	22.2	20.9	9.7
K-feldspar	0	0.8	0.1	0.7	2.6	0.2	6.0	0.2	1.7
Plagioclase	49.7	49.0	46.4	37.6	25.6	40.5	52.3	50.3	58.2
Biotite	23.2	13.7	25.5	24.6	33.2	32.2	15.5	18.1	14.4
Hornblende	24.9	18.4	21.0	35.6	22.5	14.0	2.9	9.0	14.9
Sphene			1.4		1.0				
Opaque	1.0	0.9	0.4	0.7	0.8	1.2	1.1	1.5	1.1
Other			0		0.9				
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

	Dinkey Creek Granodiorite—Continued								
	Markwood Creek		Deer Creek				Wishon Reservoir		
	Granodiorite	Inclusion	Tonalite	Inclusion			Granodiorite	Aggregate	Inclusion
	MK1-H	MK2-I	DR1a-H	DR1b-I	DR4a-I	DR4b-I	WR1-H	WR2-A	WR3-I
Quartz	23.1	1.2	13.7	1.6	0.3	2.1	23.7	14.3	1.7
K-feldspar	8.5	0.4	0	0.1	0	0	6.0	9.0	0
Plagioclase	47.5	42.9	49.0	43.0	44.4	53.7	46.4	24.3	43.0
Biotite	11.7	18.0	16.7	18.8	2.6	5.9	13.0	23.9	23.7
Hornblende	8.4	36.7	19.2	34.7	46.9*	34.7*	10.1	26.0	30.6
Sphene				0.1	0	1.7			
Opaque	0.8	0.8	1.4	1.8	4.5	3.3	0.8	0.2	1.0
Other				1.2	0.3	0.6			
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Table 2. Modal analyses, in volume percent, of mafic inclusions, aggregates, dikes, and their host granitoids and associated gabbroid—Continued

	Granodiorite of Red Lake			Mount Givens Granodiorite					
	Chinese Peak			Mono Hot Springs		Camp 61			
	Granodiorite	Inclusion		Granodiorite	Inclusion	Granodiorite		Aggregate	
	RL1-H	RL2-I	RL3-I	MHa1-H	MHa2-I	MHb1-H	MHb3-H	MHb4-A	MHb5-A
Quartz	26.8	1.7	3.7	26.2	4.2	21.2	24.7	15.7	14.4
K-feldspar	12.3	3.3	2.7	16.8	5.5	19.6	20.5	12.2	19.1
Plagioclase	49.1	56.1	53.6	39.9	52.9	44.5	39.3	32.7	35.4
Biotite	8.7	20.1	32.9	8.2	8.0	8.3	9.0	18.3	14.2
Hornblende	2.1	16.5	6.0	7.6	28.3	5.0	5.3	16.9	12.2
Sphene	}	}	}	}	}	}	}	1.8	1.1
Opaque	1.0	1.5	1.1	1.3	1.1	1.4	1.2	2.2	1.6
Other	}	}	}	}	}	}	}	0.2	1.0
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

	Mount Givens Granodiorite—Continued								
	Camp 61—Continued								
	Inclusion								
	MHb6-I	MHb7-I	MHb8-I	MHb9a-Im	MHb9b-Ic	MHb10-I	MHb11-I	MHb12-I	MHb15-I
Quartz	4.8	2.3	4.5	0.3	2.3	1.8	12.5	7.8	16.6
K-feldspar	0	0.4	10.5	0	0	0	0.4	0.4	0.8
Plagioclase	51.4	55.2	25.6	47.9	55.3	44.9	54.7	58.6	50.4
Biotite	22.3	20.9	31.7	17.6	18.2	18.3	19.3	20.9	16.7
Hornblende	20.6	19.4	26.7	32.3	22.9	33.6	10.6	10.4	13.7
Sphene	}	}	}	}	}	}	0.7	0.3	}
Opaque	0.9	1.8	1.0	1.9	1.3	1.4	1.1	1.2	1.8
Other	}	}	}	}	}	}	0.7	0.4	}
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

	Mount Givens Granodiorite—Continued						
	Courtright Reservoir						
	Aggregate			Inclusion			
	CR3a-A	CR4-A	CR5-A	CR3b-I	CR6-I	CR8-I	CR10-I
Quartz	18.0	24.0	6.4	7.2	2.7	0.6	2.1
K-feldspar	0	3.4	9.8	4.4	0.9	0	1.6
Plagioclase	35.9	34.8	28.6	42.0	53.8	50.9	38.5
Biotite	37.6	26.8	39.9	8.4	19.8	16.7	5.3
Hornblende	5.0	7.4	9.2	36.0	18.2	29.6	51.6
Sphene	1.6	1.6	3.2	}	2.5	0.7	}
Opaque	1.4	1.4	2.5	2.0	0.9	1.2	0.9
Other	0.5	0.6	0.4	}	1.2	0.3	}
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0

*Includes clinopyroxene.

Table 3. Chemical and trace-element analyses of mafic inclusions, aggregates, dikes, and their host granitoids, associated gabbroid, and aplite

[Chemical analyses by A.J. Bartel, K. Stewart, J. Taggart, H. Neiman, E. Engleman, M. Maneer, C. Stone, G. Mason, J. Ryder, and L. Espos using a combination of X-ray fluorescence and chemical methods. Trace-element analyses by P. Bruggman, J. Budahn, R. Knight, L.J. Schwartz, J. Storey, S. Danahey, R.B. Vaughn, J.G. Crock, J. Consul, and R. Lerner using X-ray fluorescence, instrumental neutron-activation, delayed neutron, induced coupled plasma, or emission-spectrographic methods. —, no data.]

	Bass Lake Tonalite									
	Hetch Hetchy									
	Tonalite	Aggregate		Inclusion	Dike rock					
	HH1-H	HH2-A	HH3-A	HH4-I	HH5a-D	HH5b-D	HH5c-D	HH5d-D	HH6a-D	HH6b-D
Chemical analyses (weight percent)										
SiO_2	66.9	57.2	59.4	50.4	51.9	53.4	56.2	57.2	53.1	54.4
Al_2O_3	15.2	17.5	17.0	19.1	18.8	18.3	17.6	17.5	18.4	17.7
Fe_2O_3	1.49	2.32	2.33	3.38	2.76	2.55	2.48	2.47	2.51	2.61
FeO	2.27	4.52	4.13	6.05	5.37	5.09	4.72	4.40	5.3	5.22
MgO	1.51	3.20	2.93	4.44	4.11	3.81	3.66	3.38	4.17	4.01
CaO	3.62	6.59	6.07	7.81	7.80	7.55	7.11	6.95	8.15	7.74
Na_2O	2.96	3.28	3.24	3.71	3.72	3.61	3.57	3.46	3.68	3.51
K_2O	3.47	2.03	1.89	2.07	1.98	2.05	1.86	1.78	1.71	1.62
H_2O+	0.38	0.73	0.72	1.03	0.94	0.94	0.88	0.85	0.93	1.00
H_2O-	0.19	0.20	0.20	0.22	0.25	0.25	0.24	0.25	0.22	0.19
TiO_2	0.50	0.91	0.85	1.11	1.43	1.33	1.19	1.08	1.18	1.18
P_2O_5	0.12	0.23	0.22	0.29	0.38	0.35	0.25	0.25	0.23	0.19
MnO	0.06	0.13	0.12	0.19	0.16	0.13	0.12	0.11	0.15	0.14
CO_2	<0.02	0.07	0.06	0.07	<0.02	<0.02	<0.02	<0.02	<0.02	<0.02
Total	98.7	98.9	99.2	99.9	99.6	99.4	99.9	99.7	99.7	99.5
Trace-element analyses (parts per million)										
Rb	103.	65.3	68.8	86.1	74.1	63.8	61.3	54.9	58.1	61.8
Sr	360.	488.	455.	486.	665.	654.	619.	595.	580.	569.
Cs	—	—	—	—	—	—	—	—	—	—
Ba	1544.	707.	591.	423.	507.	721.	648.	617.	582.	576.
La	18.4	37.4	19.0	21.9	21.7	24.5	23.9	23.5	23.7	20.9
Ce	35.9	78.3	40.1	54.1	50.8	53.6	48.8	48.3	49.3	42.8
Pr	1.0	3.5	3.5	4.8	3.6	0.6	1.7	1.5	2.0	1.0
Nd	13.9	38.4	24.0	34.6	31.5	30.0	24.7	24.6	25.3	21.4
Sm	1.3	5.4	3.8	5.9	4.0	1.5	1.7	1.5	1.8	1.3
Eu	1.00	1.82	1.51	1.96	2.14	2.11	1.63	1.60	1.79	1.51
Gd	2.5	8.0	5.6	8.1	6.0	5.9	5.0	4.9	5.4	5.0
Tb	0.4	0.9	0.7	1.0	0.6	0.7	0.7	0.6	0.5	0.5
Dy	1.6	6.9	4.6	7.1	4.0	4.1	3.6	3.5	4.2	4.1
Ho	0.34	1.35	0.91	1.38	0.71	0.69	0.66	0.68	0.79	0.77
Er	1.0	3.5	2.5	3.8	1.7	1.5	1.6	1.6	2.0	2.2
Tm	0.07	0.54	0.31	0.52	0.20	0.22	0.20	0.20	0.25	0.27
Yb	1.23	3.25	2.30	3.58	1.48	1.35	1.45	1.48	1.90	1.76
Lu	0.18	0.47	0.33	0.50	0.18	0.15	0.18	0.19	0.25	0.26
Y	12.	28.	21.	29.	18.	17.	16.	15.	21.	19.
Zr	134.	164.	119.	152.	145.	137.	131.	123.	132.	118.
Hf	—	—	—	—	—	—	—	—	—	—
Nb	3.	7.	8.	8.	7.	7.	5.	5.	6.	3.
Ta	—	—	—	—	—	—	—	—	—	—
Th	12.8	10.9	4.5	4.6	3.99	5.76	5.76	5.00	4.4	6.16
U	3.19	1.80	3.03	2.73	1.80	3.21	2.29	3.33	2.55	2.38
Sc	7.5	22.	18.	24.	18.	16.	19.	17.	23.	24.
V	57.	130.	120.	170.	180.	160.	180.	160.	200.	220.
Cr	—	—	—	—	—	—	—	—	—	—
Co	8.	19.	16.	22.	22.	22.	23.	19.	23.	25.
Zn	—	—	—	—	—	—	—	—	—	—
Sb	—	—	—	—	—	—	—	—	—	—

Table 3. Chemical and trace-element analyses of mafic inclusions, aggregates, dikes, and their host granitoids, associated gabbroid, and aplite—Continued

	Bass Lake Tonalite—Continued									
	Hetch Hetchy—Continued			North Fork						Meadow Lakes
	Dike rock		Aplite	Tonalite	Inclusion		Dike rock		Gabbro	Tonalite
	HH6c-D	HH6d-D	HH6e-AP	NF1-H	NF2-I	NF3-I	NF4a-D	NF4b-D	NF5-G	ML1-H
Chemical analyses (weight percent)										
SiO_2	54.3	58.4	72.5	63.6	51.3	51.2	49.6	48.4	58.6	62.4
Al_2O_3	18.1	17.4	14.0	16.8	17.9	18.2	20.1	19.8	14.8	16.9
Fe_2O_3	2.30	2.44	0.81	1.12	2.00	2.12	1.33	1.73	2.21	1.48
FeO	5.20	4.23	0.92	3.30	7.55	7.35	6.64	7.35	5.59	3.56
MgO	3.93	3.11	0.55	2.04	4.62	4.47	4.22	4.78	4.66	2.24
CaO	7.71	6.59	2.25	5.13	8.31	7.59	10.0	9.43	7.95	5.59
Na_2O	3.34	3.70	2.42	3.43	3.46	3.34	2.62	3.16	2.22	3.44
K_2O	1.77	1.62	4.93	1.88	1.75	2.15	1.40	1.46	1.53	1.97
H_2O+	0.96	0.78	0.27	0.79	1.17	1.30	1.15	1.30	1.03	0.73
H_2O-	0.22	0.18	0.16	0.19	0.24	0.25	0.31	0.30	0.20	0.05
TiO_2	1.12	0.98	0.23	0.53	0.77	0.85	1.71	1.26	0.60	0.86
P_2O_5	0.25	0.21	0.06	0.11	0.13	0.17	0.44	0.32	0.21	0.19
MnO	0.11	0.13	—	0.07	0.22	0.19	0.10	0.14	0.17	0.08
CO_2	<0.02	<0.02	<0.02	0.02	0.06	0.02	0.02	0.02	0.03	0.03
Total	99.3	99.8	99.1	99.0	99.5	99.2	99.6	99.5	99.8	99.5
Trace-element analyses (parts per million)										
Rb	54.4	68.8	90.1	66.5	61.7	87.8	34.3	40.6	43.9	64.
Sr	637.	525.	385.	276.	245.	253.	769.	620.	450.	513.
Cs	—	—	—	2.61	2.55	3.75	0.94	0.77	1.86	—
Ba	657.	419.	1693.	587.	336.	438.	519.	556.	588.	718.
La	25.7	24.6	35.8	11.8	12.5	15.3	17.5	15.4	14.0	—
Ce	52.3	47.4	63.4	25.3	37.0	44.2	45.4	39.1	29.7	—
Pr	1.8	2.2	4.3	2.8	5.1	5.2	5.7	4.8	3.3	—
Nd	26.6	22.6	22.8	13.0	29.5	25.4	30.2	24.4	14.7	—
Sm	1.9	2.1	2.0	3.1	8.3	6.9	6.6	5.7	3.8	—
Eu	1.65	1.54	0.90	0.93	1.48	1.34	1.87	1.72	1.13	—
Gd	5.3	4.8	3.2	3.6	9.7	7.7	5.6	5.2	3.9	—
Tb	0.6	0.7	0.2	0.6	1.6	1.3	0.8	0.9	0.6	—
Dy	3.9	3.6	2.4	3.4	10.6	8.7	4.4	5.1	4.3	—
Ho	0.72	0.73	0.45	0.78	2.46	1.90	0.89	1.07	0.95	—
Er	1.7	1.9	1.2	2.2	7.1	5.6	2.2	2.8	2.6	—
Tm	0.20	0.22	0.16	0.33	1.08	0.84	0.27	0.38	0.38	—
Yb	1.44	1.83	1.35	2.04	6.75	5.56	1.54	2.20	2.20	—
Lu	0.35	0.25	0.19	0.30	1.01	0.85	0.20	0.32	0.31	—
Y	18.	17.	15.	19.	48.	38.	17.	23.	21.	20.
Zr	128.	122.	127.	95.	106.	92.	57.	104.	91.	162.
Hf	—	—	—	3.18	3.57	3.09	1.40	2.72	1.96	—
Nb	6.	8.	6.	3.	6.	7.	9.	6.	6.	10.
Ta	—	—	—	0.576	0.578	0.591	0.554	0.394	0.324	—
Th	4.22	7.94	17.9	5.47	3.36	6.96	2.12	0.96	2.98	—
U	0.78	4.78	4.00	3.08	0.879	1.08	0.681	0.545	0.969	—
Sc	17.	18.	3.	13.8	38.2	31.5	19.0	26.1	33.3	—
V	160.	170.	20.	—	—	—	—	—	—	—
Cr	—	—	—	16.2	57.2	32.3	11.0	22.1	88.5	—
Co	20.	20.	3.	12.1	24.3	24.9	22.8	25.0	25.5	—
Zn	—	—	—	—	—	—	—	—	—	—
Sb	—	—	—	0.169	—	—	—	0.084	0.16	—

Table 3. Chemical and trace-element analyses of mafic inclusions, aggregates, dikes, and their host granitoids, associated gabbroid, and aplite—Continued

	Bass Lake Tonalite—Continued		Dinkey Creek Granodiorite								
	Meadow Lakes—Continued	Stevenson Creek	Shaver Lake quarry								
	Inclusion	Dike rock	Granodiorite		Inclusion						
	ML2-I	SC1-D	SQ1-H	SQ2-H	SQ3-I	SQ4-I	SQ5-I	SQ6-I	SQ7-I	SQ8-I	SQ9-I
Chemical analyses (weight percent)											
SiO_2	48.7	54.4	66.6	65.1	52.2	63.2	51.3	57.1	54.6	58.3	56.6
Al_2O_3	18.5	18.3	15.1	15.3	17.8	15.6	17.8	17.1	17.4	15.7	17.6
Fe_2O_3	2.91	1.87	1.16	1.53	2.54	1.74	2.98	2.35	2.42	2.31	2.62
FeO	7.37	5.78	2.82	3.12	6.71	4.14	7.22	5.38	5.99	5.63	5.67
MgO	4.51	3.95	1.58	1.76	4.16	2.77	4.26	3.25	3.85	3.62	2.87
CaO	8.56	7.55	3.87	4.49	7.81	5.65	7.50	6.80	7.42	6.46	5.78
Na_2O	3.61	3.19	2.70	2.96	3.51	3.18	3.52	3.44	3.46	3.11	3.62
K_2O-	1.64	1.66	4.03	3.04	1.91	1.88	2.21	1.86	1.82	1.88	2.36
H_2O+	1.20	0.97	0.58	0.60	1.12	0.72	1.14	0.86	0.88	0.81	0.88
H_2O	0.06	0.18	0.04	0.16	0.05	0.22	0.28	0.20	0.24	0.22	0.20
TiO_2	1.99	1.12	0.55	0.64	1.12	0.82	1.16	1.01	1.05	0.96	1.28
P_2O_5	0.39	0.22	0.12	0.14	0.23	0.17	0.21	0.22	0.21	0.20	0.30
MnO	0.21	0.15	0.06	0.08	0.18	0.10	0.21	0.14	0.16	0.15	0.12
CO_2	0.05	0.07	0.04	<0.02	0.06	<0.02	0.08	0.13	0.07	0.08	0.08
Total	99.7	99.4	99.3	98.9	99.4	100.2	99.9	99.8	99.6	99.4	100.0
Trace-element analyses (parts per million)											
Rb	56.	64.	132.	125.	103.	101.	142.	110.	96.4	161.	110.
Sr	483.	455.	310.	330.	336.	328.	292.	321.	344.	321.	281.
Cs	—	—	—	—	—	—	—	—	—	—	—
Ba	460.	457.	954.	755.	324.	323.	358.	359.	336.	321.	299.
La	—	14.0	—	21.2	—	13.9	18.5	15.2	18.9	26.8	12.7
Ce	—	29.5	—	45.7	—	34.9	54.1	41.2	53.7	57.0	35.8
Pr	—	2.0	—	5.0	—	3.9	7.9	3.9	6.4	5.9	5.2
Nd	—	17.9	—	21.9	—	19.7	39.0	29.4	36.3	26.7	25.0
Sm	—	1.8	—	3.9	—	3.5	8.1	5.3	6.6	4.8	5.6
Eu	—	1.49	—	1.18	—	1.13	1.70	1.46	1.70	1.19	1.19
Gd	—	4.9	—	4.6	—	4.5	10.2	7.4	7.9	5.5	6.4
Tb	—	0.4	—	0.6	—	0.6	2.0	1.1	1.3	0.7	1.0
Dy	—	3.8	—	3.4	—	2.3	9.3	6.5	7.0	3.9	5.2
Ho	—	0.68	—	0.74	—	0.64	1.85	1.28	1.38	0.77	1.01
Er	—	1.7	—	1.9	—	1.7	5.7	3.6	3.8	1.9	3.0
Tm	—	0.17	—	0.20	—	0.14	0.74	0.48	0.52	0.27	0.42
Yb	—	1.59	—	1.96	—	1.73	4.77	3.33	3.72	1.78	2.97
Lu	—	0.19	—	0.29	—	0.26	0.79	0.49	0.54	0.25	0.41
Y	31.	17.	21.	22.	40.	21.	37.	27.	31.	21.	28.
Zr	167.	109.	116.	143.	180.	137.	152.	134.	152.	176.	130.
Hf	—	—	—	—	—	—	—	—	—	—	—
Nb	19.	5.	8.	9.	14.	10.	16.	10.	13.	11.	13.
Ta	—	—	—	—	—	—	—	—	—	—	—
Th	—	—	—	6.5	—	6.41	4.5	—	4.1	10.4	4.5
U	—	1.7	—	4.18	—	3.21	3.84	4.79	1.51	2.17	2.97
Sc	—	19.	—	10.	—	14.	30.	19.	22.	18.	18.
V	—	140.	—	80.	—	108.	190.	130.	170.	145.	125.
Cr	—	—	—	—	—	—	—	—	—	—	—
Co	—	21.	—	12.	—	16.	21.	17.	22.	17.	17.
Zn	—	—	—	—	—	—	—	—	—	—	—
Sb	—	—	—	—	—	—	—	—	—	—	—

Table 3. Chemical and trace-element analyses of mafic inclusions, aggregates, dikes, and their host granitoids, associated gabbroid, and aplite—Continued

	Dinkey Creek Granodiorite—Continued									
	Shaver Lake quarry—Continued				Camp Sierra			Markwood Creek		Deer Creek
	Inclusion	Dike rock			Grano-diorite	Inclusion		Grano-diorite	Inclusion	Grano-diorite
	SQ10- I	SQ12-aD	SQ12-Dx	SQ12-Di	CS1-H	CS2-I	CS3-I	MK1-H	MK2-I	DR1-aH
	Chemical analyses (weight percent)									
SiO_2	50.6	53.9	52.6	54.4	62.4	60.5	59.2	63.3	50.1	57.2
Al_2O_3	18.7	14.1	17.6	13.1	16.2	16.8	17.6	16.2	17.4	17.3
Fe_2O_3	2.90	3.51	2.81	3.37	1.46	2.12	1.79	1.76	3.19	2.27
FeO	6.84	8.18	7.19	8.94	4.34	4.80	4.52	3.47	7.66	5.09
MgO	4.03	4.61	3.98	4.81	2.30	2.15	2.41	2.16	4.83	3.57
CaO	7.14	6.28	6.88	5.73	5.22	5.46	6.19	5.06	8.32	7.26
Na_2O	3.63	2.57	3.65	2.21	2.98	3.50	3.52	3.19	3.35	2.98
K_2O	2.52	2.97	2.39	3.32	2.58	1.93	1.87	2.59	1.86	1.73
H_2O+	1.14	1.16	0.93	1.18	0.73	0.69	0.79	0.79	1.20	0.82
H_2O-	0.23	0.24	0.20	0.20	0.13	0.11	0.16	0.10	0.14	0.15
TiO_2	1.27	1.39	1.25	1.57	0.80	1.00	0.92	0.73	1.16	1.03
P_2O_5	0.29	0.30	0.27	0.33	0.17	0.31	0.20	0.16	0.23	0.23
MnO	0.18	0.24	0.20	0.24	0.09	0.08	0.09	0.09	0.22	0.13
CO_2	0.11	<0.02	<0.02	0.07	0.01	0.01	0.01	0.07	0.09	0.14
Total	99.6	99.5	100.0	99.5	99.4	99.5	99.3	99.7	99.8	99.9
	Trace-element analyses (parts per million)									
Rb	169.	161.	156.	196.	105.	96.	86.	100.	91.	62.6
Sr	330.	207.	276.	180.	378.	424.	431.	369.	318..	471.
Cs	—	—	—	—	—	—	—	—	—	—
Ba	237.	443.	316.	542.	760.	506.	573.	764.	255.	516.
La	—	74.1	26.1	170.0	—	—	—	—	28.9	16.9
Ce	—	155.0	64.7	322.0	—	—	—	—	60.3	41.0
Pr	—	18.8	9.7	33.1	—	—	—	—	—	7.7
Nd	—	73.7	40.8	118.0	—	—	—	—	31.6	26.9
Sm	—	15.0	9.9	18.3	—	—	—	—	—	5.1
Eu	—	2.03	1.66	2.34	—	—	—	—	1.71	1.76
Gd	—	14.3	9.9	17.0	—	—	—	—	—	7.1
Tb	—	2.2	1.6	2.5	—	—	—	—	—	1.1
Dy	—	12.8	9.1	12.6	—	—	—	—	—	5.2
Ho	—	2.46	1.79	2.65	—	—	—	—	1.01	1.17
Er	—	6.8	5.1	7.0	—	—	—	—	—	2.8
Tm	—	0.99	0.73	0.90	—	—	—	—	0.30	0.44
Yb	—	6.49	4.73	6.92	—	—	—	—	2.55	2.60
Lu	—	0.91	0.71	0.99	—	—	—	—	0.41	0.36
Y	33.	56.	40.	57.	27.	21.	23.	28.	47.	23.
Zr	177.	266.	184.	304.	191.	145.	162.	146.	146.	136.
Hf	—	—	—	—	—	—	—	—	—	—
Nb	16.	22.	19.	25.	9.	8.	7.	10.	15.	9.
Ta	—	—	—	—	—	—	—	—	—	—
Th	—	41.5	11.2	67.5	—	—	—	—	7.47	3.00
U	—	7.44	4.49	8.33	—	—	—	—	3.22	1.20
Sc	—	40.	30.	40.	—	—	—	—	22.	48.
V	—	210.	180.	225.	—	—	—	—	160.	300.
Cr	—	—	—	—	—	—	—	—	—	—
Co	—	22.	20.	24.	—	—	—	—	18.	25.
Zn	—	—	—	—	—	—	—	—	—	—
Sb	—	—	—	—	—	—	—	—	—	—

Table 3. Chemical and trace-element analyses of mafic inclusions, aggregates, dikes, and their host granitoids, associated gabbroid, and aplite—Continued

	Dinkey Creek Granodiorite–Continued							Granodiorite of Red Lake		
	Deer Creek–Continued				Wishon Reservoir			Chinese Peak		
	Aggregate	Inclusion			Grano-diorite	Aggregate	Inclusion	Grano-diorite	Inclusion	
	DR2-S	DR3-I	DR4a-I	DR5-I	WR1-H	WR2-A	WR3-I	RL1-H	RL2-I	RL3-I
	Chemical analyses (weight percent)									
SiO_2	55.6	48.3	46.1	49.1	60.8	56.0	48.1	69.3	52.7	53.0
Al_2O_3	15.5	18.4	19.9	20.0	16.0	13.4	17.8	15.2	18.2	18.6
Fe_2O_3	3.00	3.42	3.52	1.81	1.46	3.20	2.82	1.16	2.67	2.54
FeO	6.57	6.82	7.63	6.74	4.64	7.29	8.44	1.58	5.78	4.91
MgO	4.94	5.29	5.18	6.77	2.80	5.03	5.06	0.94	4.10	3.51
CaO	7.14	10.20	10.80	10.90	6.07	6.40	8.38	3.20	5.99	6.81
Na_2O	2.46	1.90	2.42	2.23	2.92	2.02	3.02	3.58	4.21	4.29
K_2O-	2.03	1.27	0.46	0.51	2.25	3.14	2.47	2.90	2.62	2.11
H_2O+	1.29	1.98	1.54	1.15	0.77	1.38	1.38	0.37	1.12	0.99
H_2O	0.31	0.31	0.22	0.19	0.05	0.26	0.10	0.04	0.19	0.24
TiO_2	1.32	1.50	1.60	0.57	0.88	1.36	1.35	0.42	1.12	1.30
P_2O_5	0.29	0.28	0.31	0.12	0.19	0.28	0.44	0.14	0.31	0.42
MnO	0.17	0.15	0.19	0.18	0.10	0.21	0.20	0.04	0.20	0.19
CO_2	<0.02	0.32	0.25	<0.02	0.03	0.09	0.07	0.04	0.06	<0.02
Total	100.6	100.1	100.1	100.3	99.0	100.1	99.6	98.9	99.3	98.9
	Trace-element analyses (parts per million)									
Rb	50.6	76.6	15.4	13.1	96.1	124.	1240.	93.	125.	95.
Sr	471.	382.	578.	575.	402.	257.	380.	572.	552.	539.
Cs	—	—	—	4.23	—	7.4	—	1.69	8.24	—
Ba	290.	550.	158.	186.	573.	732.	431.	1156.	1016.	556.
La	31.3	18.5	14.6	30.1	76.0	23.8	—	29.4	15.8	31.9
Ce	71.8	44.4	34.0	57.2	144.0	61.1	—	52.0	29.4	68.0
Pr	5.2	10.0	6.0	4.9	—	13.2	—	5.5	3.8	5.4
Nd	39.8	28.2	19.9	27.0	64.7	40.0	—	23.0	21.0	32.0
Sm	6.7	8.5	5.8	4.3	5.7	9.7	8.4	3.9	3.7	3.8
Eu	1.64	1.85	1.59	1.17	2.23	1.52	—	0.82	0.81	2.38
Gd	7.3	8.1	6.4	4.2	4.8	12.6	8.0	3.6	4.0	5.5
Tb	1.2	1.1	1.0	0.7	0.7	2.0	1.2	0.3	0.4	0.8
Dy	6.4	7.0	5.6	3.5	—	10.5	—	1.7	2.4	3.4
Ho	1.33	1.10	0.68	—	2.03	—	—	0.35	0.52	0.61
Er	3.2	3.7	3.1	1.9	—	5.5	—	0.9	1.4	1.4
Tm	0.54	0.45	0.26	0.32	0.69	0.47	—	0.12	—	0.19
Yb	3.56	2.89	1.76	2.22	5.21	3.52	—	0.91	1.22	1.45
Lu	0.50	0.44	0.29	0.36	0.73	0.52	—	0.13	0.21	0.22
Y	26.	31.	27.	18.	26.	38.	37.	14.	17.	17.
Zr	77.	149.	51.	92.	143.	185.	162.	162.	166.	156.
Hf	—	—	—	4.95	—	4.71	—	3.98	4.58	—
Nb	9.	11.	11.	5.	13.	14.	12.	8.	10.	10.
Ta	—	—	—	1.48	—	1.07	—	0.65	0.59	—
Th	5.56	—	—	17.8	17.9	3.58	—	10.9	3.31	8.05
U	2.37	0.367	0.443	6.2	7.68	2.6	—	1.92	3.3	3.14
Sc	29.	34.	14.	16.6	32.	29.6	—	3.53	21.1	13.
V	200.	250.	85.	—	7.68	—	—	—	—	160.
Cr	—	—	—	15.8	—	87.1	—	3.4	19.9	—
Co	24.	26.	29.	14.8	25.	31.7	—	4.13	19.2	18.
Zn	—	—	—	82.	—	163.	—	56.	212.	—
Sb	—	—	—	0.45	—	0.58	—	0.20	1.51	—

Table 3. Chemical and trace-element analyses of mafic inclusions, aggregates, dikes, and their host granitoids, associated gabbroid, and aplite—Continued

	Mount Givens Granodiorite									
	Camp 61									
	Grano-diorite	Inclusion	Granodiorite			Aggregate		Inclusion		
	MHa1-H	MHa2-I	MHb1-H	MHb2-H	MHb3-H	MHb4-A	MHb5-A	MHb6-I	MHb7-I	MHb8-I
Chemical analyses (weight percent)										
SiO_2	64.7	52.3	66.9	67.1	66.6	54.9	61.7	55.3	53.4	52.0
Al_2O_3	15.9	17.6	15.4	15.2	15.1	13.3	15.4	17.0	17.6	17.1
Fe_2O_3	1.58	3.12	1.54	1.71	1.72	4.74	2.61	2.08	3.70	2.47
FeO	2.52	5.95	2.13	2.01	2.15	6.53	3.49	5.51	5.18	5.67
MgO	1.74	4.55	1.55	1.53	1.60	5.08	2.76	4.63	3.98	5.48
CaO	4.60	7.91	3.78	3.76	3.96	4.80	4.66	6.69	6.90	5.72
Na_2O	3.25	3.91	3.26	3.18	3.36	2.24	3.26	3.90	4.47	3.63
K_2O	2.79	1.61	3.60	3.56	3.08	3.93	3.31	2.00	1.42	3.93
H_2O+	0.48	0.95	0.43	0.53	0.51	1.26	0.72	0.83	1.26	1.21
H_2O	0.04	0.07	0.03	0.14	0.25	0.18	0.16	0.03	0.06	0.08
TiO_2	0.54	0.96	0.57	0.53	0.62	1.38	0.78	0.99	0.80	0.99
P_2O_5	0.14	0.20	0.16	0.16	0.16	0.35	0.22	0.26	0.25	0.42
MnO	0.07	0.23	0.06	0.06	0.06	0.23	0.13	0.21	0.21	0.21
CO_2	0.04	0.04	0.04	<0.02	<0.02	<0.02	<0.02	0.03	0.09	0.05
Total	98.4	99.4	99.5	99.5	99.2	98.9	99.2	99.5	99.3	99.0
Trace-element analyses (parts per million)										
Rb	116.	94.	130.	129.	126.	207.	142.	141.	77.	291.
Sr	431.	376.	414.	425.	434.	262.	387.	356.	403.	360.
Cs	—	—	6.79	—	—	—	—	13.8	4.04	19.8
Ba	616.	195.	764.	863.	732.	845.	630.	359.	281.	611.
La	—	—	28.4	35.2	54.3	64.7	45.9	25.2	25.0	33.9
Ce	—	—	54.5	60.2	93.0	123.0	86.2	58.6	56.0	61.2
Pr	—	—	5.8	3.9	8.1	13.6	8.5	6.3	7.0	6.5
Nd	—	—	25.0	23.0	34.1	56.6	39.2	27.0	37.0	28.0
Sm	—	—	4.6	1.6	3.8	10.1	6.4	7.2	8.7	5.9
Eu	—	—	0.88	1.05	1.29	1.69	1.44	1.08	1.27	0.80
Gd	—	—	4.2	4.1	5.5	10.3	7.3	6.4	6.4	5.3
Tb	—	—	0.4	0.4	0.6	1.5	1.1	0.9	1.2	0.8
Dy	—	—	2.7	2.7	3.3	8.2	5.7	5.1	6.7	4.5
Ho	—	—	0.60	0.56	0.68	1.56	1.09	0.98	1.33	0.95
Er	—	—	1.7	1.4	1.7	4.1	2.9	2.2	3.2	2.5
Tm	—	—	0.24	0.14	0.14	0.56	0.37	0.25	0.39	0.33
Yb	—	—	1.52	1.49	1.81	4.06	2.86	2.35	2.76	2.26
Lu	—	—	0.25	0.21	0.26	0.61	0.42	0.39	0.42	0.35
Y	23.	42.	22.	21.	18.	38.	27.	31.	32.	31.
Zr	123.	141.	136.	130.	129.	266.	184.	151.	111.	225.
Hf	—	—	4.52	—	—	—	—	4.68	3.84	5.19
Nb	5.	15.	10.	8.	7.	20.	12.	11.	14.	8.
Ta	—	—	1.31	—	—	—	—	1.09	1.15	0.96
Th	—	—	24.1	22.3	21.1	38.2	27.6	8.25	8.77	7.83
U	—	—	5.6	5.00	4.30	10.20	9.20	8.1	15.8	4.1
Sc	—	—	7.82	7.5	6.5	30.	20.	28.8	35.9	22.4
V	—	—	—	66.	64.	190.	100.	—	—	—
Cr	—	—	9.4	—	—	—	83.2	30.2	—	92.9
Co	—	—	8.14	9.	9.	23.	13.	25.8	23.0	28.2
Zn	—	—	57.	—	—	—	161.	165.	—	188.
Sb	—	—	0.67	—	—	—	—	0.44	1.42	0.50

Table 3. Chemical and trace-element analyses of mafic inclusions, aggregates, dikes, and their host granitoids, associated gabbroid, and aplite—Continued

	Mount Givens Granodiorite—Continued									
	Camp 61—Continued								Courtright Reservoir	
	Inclusion—Continued								Grano-diorite	Aggregate
	MHb9a-Im	MHb9b-Ic	MHb10-I	MHb11-I	MHb12-I	MHb13-I	MHb14-I	MHb15-I	CR1-H	CR2a-A
	Chemical analyses (weight percent)									
SiO_2	50.6	54.5	50.8	56.7	55.3	52.2	50.4	59.2	65.5	61.2
Al_2O_3	17.8	17.2	17.7	17.1	17.6	17.1	20.0	17.3	17.0	15.5
Fe_2O_3	2.89	2.40	2.86	2.56	3.26	3.79	3.45	2.10	0.80	2.04
FeO	6.40	5.22	6.13	4.76	4.48	5.24	5.54	3.97	2.01	3.95
MgO	5.33	4.65	5.42	3.86	3.65	5.55	3.77	3.08	0.95	2.70
CaO	6.87	7.23	7.41	5.39	7.03	6.60	6.55	5.93	2.73	4.07
Na_2O	3.89	4.07	3.95	3.79	4.19	3.98	4.74	4.09	3.39	3.00
K_2O-	2.53	1.67	2.18	2.23	1.35	2.25	2.26	1.49	5.95	4.02
H_2O+	1.27	1.05	1.20	1.13	0.97	1.24	1.07	0.78	0.43	0.80
H_2O	0.30	0.19	0.18	0.18	0.26	0.21	0.19	0.21	0.17	0.19
TiO_2	1.23	1.10	1.24	0.96	1.00	1.07	1.25	0.82	0.47	1.13
P_2O_5	0.37	0.32	0.37	0.24	0.25	0.27	0.47	0.20	0.14	0.33
MnO	0.23	0.20	0.23	0.16	0.17	0.25	0.16	0.13	0.06	0.10
CO_2	<0.02	<0.02	<0.02	<0.02	<0.02	<0.02	<0.02	<0.02	<0.02	0.06
Total	99.7	99.8	99.7	99.1	99.5	99.8	99.9	99.3	99.6	99.1
	Trace-element analyses (parts per million)									
Rb	167.	103.	172.	177.	78.8	210.	193.	98.1	184.	120.
Sr	362.	364.	375.	419.	426.	329.	410.	403.	484.	533.
Cs	—	—	—	—	—	—	—	—	—	—
Ba	484.	370.	355.	379.	362.	377.	293.	293.	3161.	2622.
La	35.8	40.6	40.0	21.1	42.0	27.3	49.6	25.6	36.6	56.6
Ce	70.3	84.5	89.6	37.5	85.2	55.2	89.5	50.2	57.9	122.0
Pr	5.5	5.2	8.3	1.2	4.4	7.0	5.8	3.8	4.8	12.5
Nd	36.7	38.6	47.4	16.3	30.2	28.4	33.9	25.2	18.4	52.4
Sm	6.0	4.5	7.4	1.6	1.0	6.1	3.0	3.2	1.7	7.5
Eu	1.29	1.43	1.67	0.85	1.48	1.00	1.29	1.02	0.94	2.12
Gd	8.9	8.3	10.2	4.3	5.6	6.0	5.7	5.3	3.3	7.0
Tb	1.1	0.9	1.2	0.6	0.6	0.8	0.8	0.7	0.2	1.0
Dy	6.1	6.1	7.6	2.8	4.1	4.4	4.2	4.2	0.8	4.6
Ho	1.15	1.17	1.41	0.54	0.81	0.84	0.79	0.75	0.15	0.83
Er	2.9	3.0	3.7	1.3	2.0	2.2	1.9	1.9	0.4	2.0
Tm	0.33	0.40	0.47	0.11	0.29	0.28	0.23	0.21	<0.05	0.25
Yb	2.38	2.93	3.32	1.31	2.32	1.92	1.89	1.67	0.38	1.87
Lu	0.33	0.41	0.45	0.18	0.38	0.36	0.30	0.28	0.04	0.26
Y	30.	28.	33.	18.	22.	23.	21.	21.	12.	22.
Zr	160.	145.	140.	73.	154.	110.	173.	131.	130.	161.
Hf	—	—	—	—	—	—	—	—	—	—
Nb	12.	13.	13.	6.	16.	10.	10.	6.	4.	14.
Ta	—	—	—	—	—	—	—	—	—	—
Th	5.70	6.34	5.89	—	—	—	7.22	6.35	12.9	19.0
U	3.26	4.23	3.33	6.13	9.99	5.74	6.39	2.74	2.55	3.05
Sc	28.	20.	19.	20.	25.	26.	17.	16.	7.5	13.
V	170.	164.	180.	118.	136.	160.	190.	110.	45.	110.
Cr	—	—	—	—	—	—	—	—	—	—
Co	25.	21.	26.	19.	18.	24.	23.	15.	7.	12.
Zn	—	—	—	—	—	—	—	—	—	—
Sb	—	—	—	—	—	—	—	—	—	—

Table 3. Chemical and trace-element analyses of mafic inclusions, aggregates, dikes, and their host granitoids, associated gabbroid, and aplite—Continued

	Mount Givens Granodiorite—Continued									
	Courtright Reservoir—Continued									
	Aggregate—Continued			Inclusion						Hornfels
	CR3a-A	CR4-A	CR5-A	CR2b-I	CR3b-I	CR6-I	CR7-I	CR8-I	CR9-I	CR12-I
	Chemical analyses (weight percent)									
SiO_2	56.6	57.8	51.9	55.0	56.8	53.5	52.3	51.4	54.2	54.6
Al_2O_3	14.2	14.0	14.1	18.4	13.6	18.1	18.6	17.8	18.0	16.5
Fe_2O_3	2.81	2.99	5.23	2.78	2.93	3.64	3.49	3.25	2.72	1.05
FeO	6.83	5.91	6.90	5.19	6.54	4.82	4.96	5.42	5.11	5.98
MgO	4.42	3.84	5.28	3.44	5.30	4.12	4.15	5.13	4.18	2.57
CaO	4.02	4.36	4.42	5.74	6.00	6.17	6.81	7.73	5.09	16.30
Na_2O	2.48	2.55	2.25	4.21	2.81	4.29	4.57	4.42	4.24	0.94
K_2O	3.42	3.66	5.04	2.16	2.05	2.10	2.06	1.70	2.67	0.14
H_2O+	1.50	1.19	1.39	1.05	1.66	1.16	0.92	0.99	1.76	0.38
H_2O	0.25	0.30	0.26	0.25	0.23	0.29	0.21	0.24	0.29	0.15
TiO_2	2.05	1.78	2.37	1.12	0.94	1.12	1.19	1.20	1.08	1.03
P_2O_5	0.55	0.54	0.71	0.37	0.11	0.34	0.42	0.34	0.31	0.33
MnO	0.16	0.15	0.20	0.14	0.25	0.15	0.15	0.18	0.18	0.35
CO_2	<0.02	0.09	<0.02	0.07	0.07	0.06	0.06	<0.02	0.09	<0.02
Total	99.3	99.2	100.1	99.9	99.3	99.9	99.9	99.8	99.9	100.3
	Trace-element analyses (parts per million)									
Rb	163.	141.	286.	104.	84.8	115.	115.	101.	175.	3.
Sr	390.	395.	330.	553.	352.	522.	516.	510.	512.	275.
Cs	—	—	—	—	—	—	—	—	—	—
Ba	1382.	1945.	1109.	739.	819.	332.	347.	249.	403.	66.
La	79.0	83.2	96.5	28.3	23.5	38.7	41.9	41.0	31.8	30.6
Ce	192.0	188.0	210.0	58.3	53.5	69.6	79.3	85.1	56.6	56.3
Pr	21.4	21.8	24.6	5.3	6.9	5.4	4.4	7.4	5.3	6.0
Nd	82.6	85.9	91.9	26.7	31.0	26.0	34.3	32.9	19.0	26.5
Sm	12.5	13.3	14.3	3.5	6.1	2.9	3.0	5.0	3.4	5.4
Eu	3.12	3.29	3.64	1.31	1.68	1.19	1.42	1.57	0.90	1.75
Gd	10.6	12.5	11.6	4.5	6.4	4.9	5.7	5.9	3.5	6.8
Tb	1.4	1.6	1.3	0.5	0.9	0.4	0.9	0.7	0.3	1.0
Dy	6.3	8.2	7.4	2.8	4.1	2.4	3.7	3.5	2.1	6.3
Ho	1.15	1.50	1.36	0.49	0.82	0.51	0.61	0.66	0.40	1.36
Er	2.6	3.9	3.2	1.2	2.1	1.1	1.4	1.7	1.0	4.0
Tm	0.22	0.45	0.37	0.13	0.26	0.05	0.17	0.20	0.15	0.60
Yb	2.48	3.01	3.20	1.16	2.13	1.13	1.39	1.47	1.01	4.07
Lu	0.32	0.48	0.46	0.16	0.33	0.19	0.18	0.21	0.15	0.56
Y	30.	33.	29.	16.	21.	14.	20.	17.	18.	34.
Zr	289.	254.	312.	135.	101.	134.	193.	141.	136.	169.
Hf	—	—	—	—	—	—	—	—	—	—
Nb	23.	25.	21.	8.	7.	8.	12.	8.	9.	6.
Ta	—	—	—	—	—	—	—	—	—	—
Th	24.0	24.5	17.5	7.44	7.17	7.40	7.41	6.28	7.75	7.61
U	5.80	4.36	9.72	3.07	2.56	3.28	3.80	2.45	4.96	4.04
Sc	17.	18.	16.	13.	34.	16.	17.	23.	20.	27.
V	250.	220.	215.	170.	175.	180.	165.	190.	185.	260.
Cr	—	—	—	—	—	—	—	—	—	—
Co	25.	23.	27.	16.	20.	20.	19.	24.	24.	17.
Zn	—	—	—	—	—	—	—	—	—	—
Sb	—	—	—	—	—	—	—	—	—	—

Table 4. CIPW norms of mafic inclusions, aggregates, dikes, and their host granitoids, associated gabbroid, and aplite

[tr, trace; —, not calculated]

	Bass Lake Tonalite											
	Hetch Hetchy											
	Tonalite	Aggregate		Inclusion	Dike rock							
	HH1-H	HH2-A	HH3-A	HH4-I	HH5a-D	HH5b-D	HH5c-D	HH5d-D	HH6a-D	HH6b-D	HH6c-D	HH6d-D
Q	25.7	10.8	15.1	—	tr	2.9	7.7	10.3	1.6	5.2	5.3	11.7
or	20.5	12.0	11.2	12.2	11.7	12.1	11.0	10.5	10.1	9.6	10.5	9.6
ab	25.1	27.8	27.4	31.4	31.5	30.6	30.2	29.3	31.1	29.7	28.3	31.3
an	17.2	27.0	26.3	29.4	28.8	27.7	26.5	27.0	28.6	27.8	29.2	26.1
C	0.3	—	—	—	—	—	—	—	—	—	—	—
di	—	3.0	1.7	5.8	6.1	6.1	5.8	4.8	8.4	7.6	6.1	4.3
hy	6.0	11.6	11.0	3.9	12.8	11.8	11.2	10.7	12.3	12.0	12.8	10.1
mt	2.2	3.4	3.4	4.9	4.0	3.7	3.6	3.6	3.6	3.8	3.3	3.5
ne	—	—	—	—	—	—	—	—	—	—	—	—
ol	—	—	—	8.1	—	—	—	—	—	—	—	—
il	1.0	1.7	1.6	2.1	2.7	2.5	2.3	2.1	2.2	2.2	2.1	1.9
ap	0.3	0.5	0.5	0.7	0.9	0.8	0.6	0.6	0.5	0.4	0.6	0.5
cc	—	0.2	0.1	0.2	—	—	—	—	—	—	—	—
Total	98.3	98.0	98.3	98.7	98.5	93.2	98.9	98.9	98.4	98.3	98.2	99.0

	Bass Lake Tonalite—Continued									Dinkey Creek Granodiorite		
	Hetch Hetchy—Continued	North Fork						Meadow Lakes		Shaver Lake quarry		
	Aplite	Tonalite	Inclusion		Dike rock		Gabbro	Tonalite	Inclusion	Granodiorite		Inclusion
	HH6e-AP	NF1-H	NF2-I	NF3-I	NF4a-D	NF4b-D	NF5-G	ML1-H	ML2-I	SQ1-H	SQ2-H	SQ3-I
Q	33.8	20.8	—	—	0.4	—	15.6	18.6	—	23.9	23.2	0.5
or	29.1	11.1	10.3	12.7	8.3	8.6	9.0	11.6	9.7	23.8	18.0	11.3
ab	20.5	29.0	29.3	28.3	22.2	26.7	18.8	29.1	30.6	22.9	25.1	29.7
an	10.8	24.6	28.1	28.3	39.0	35.5	25.9	24.9	29.4	17.2	19.5	27.2
C	0.7	0.1	—	—	—	—	—	—	—	—	—	—
di	—	—	9.8	6.7	6.3	7.4	9.8	1.2	8.3	0.8	1.5	8.0
hy	2.0	9.5	9.3	11.1	15.9	1.0	14.6	9.0	0.7	7.0	7.2	15.1
mt	1.2	1.6	2.9	3.1	1.9	2.5	3.2	2.2	4.2	1.7	2.2	3.7
ne	—	—	—	—	—	—	—	—	—	—	—	—
ol	—	—	6.4	5.5	—	12.9	—	—	10.8	—	—	—
il	0.4	1.0	1.5	1.6	3.3	2.4	1.1	1.6	3.8	1.0	1.2	2.1
ap	0.1	0.3	0.3	0.4	1.0	0.7	0.5	0.4	0.9	0.3	0.3	0.5
cc	—	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	—	0.1
Total	98.6	98.1	98.0	97.8	98.4	97.8	98.6	98.7	98.5	98.7	98.2	98.2

	Dinkey Creek Granodiorite—Continued											
	Shaver Lake quarry—Continued										Stevenson Creek	Camp Sierra
	Inclusion—Continued							Dike rock				Granodiorite
	SQ4-I	SQ5-I	SQ6-I	SQ7-I	SQ8-I	SQ9-I	SQ10-I	SQ12a-D	SQ12a-Di	SQ12a-Dx	SC1-D	CS1-H
Q	20.4	—	9.8	5.0	12.8	8.3	—	5.8	—	7.4	5.9	18.9
or	11.1	13.1	11.0	10.8	11.1	14.0	14.9	17.6	14.1	19.6	9.8	15.3
ab	26.9	29.8	29.1	29.3	26.3	30.6	30.7	21.8	30.9	18.7	27.0	25.2
an	22.7	26.2	25.7	26.6	23.3	24.8	27.3	18.2	24.6	16.0	30.7	23.2
C	—	—	—	—	—	—	—	—	—	—	—	—
di	3.4	7.5	4.7	6.9	5.7	1.2	4.6	9.1	6.4	8.1	4.0	1.3
hy	10.2	10.5	12.3	13.8	13.4	12.9	5.0	17.3	15.2	19.5	15.4	10.7
mt	2.5	4.3	3.4	3.5	3.4	3.8	4.2	5.1	4.1	4.9	2.7	2.1
ne	—	—	—	—	—	—	—	—	—	—	—	—
ol	—	4.2	—	—	—	—	8.2	—	0.5	—	—	—
il	1.6	2.2	1.9	2.0	1.8	2.4	2.4	2.6	2.4	3.0	2.1	1.5
ap	0.4	0.5	0.5	0.5	0.5	0.7	0.7	0.7	0.6	0.8	0.5	0.4
cc	—	0.2	0.3	0.2	0.2	0.2	0.3	—	—	0.2	0.2	tr
Total	99.2	98.5	98.7	98.6	98.5	98.9	98.3	98.2	98.8	98.2	98.3	98.6

Table 4. CIPW norms of mafic inclusions, aggregates, dikes, and their host granitoids, associated gabbroid, and aplite—Continued

	Dinkey Creek Granodiorite—Continued											
	Camp Sierra—Continued		Markwood Creek		Deer Creek					Wishon Reservoir		
	Inclusion		Granodiorite	Inclusion	Granodiorite	Aggregate	Inclusion			Granodiorite	Aggregate	Inclusion
	CS2-I	CS3-I	MK1-H	MK2-I	DR1-H	DR2-A	DR3-I	DR4-I	DR5-I	WR1-H	WR2-A	WR3-I
Q	16.3	13.4	19.9	—	11.7	10.0	2.8	—	—	16.9	10.4	—
or	11.4	11.1	15.3	11.0	10.2	12.0	7.5	2.7	3.0	13.3	18.6	14.6
ab	29.6	29.8	27.0	28.4	25.2	20.8	16.1	20.5	18.9	24.7	17.1	24.1
an	24.4	26.7	22.2	27.0	28.7	25.3	37.9	42.1	43.1	23.9	18.2	27.7
C	—	—	—	—	—	—	—	—	—	—	—	—
di	0.5	2.2	1.1	10.1	4.0	6.7	7.2	6.5	8.3	3.9	9.1	8.7
hy	10.7	10.4	8.7	7.6	12.9	16.7	17.2	13.6	17.9	11.1	17.0	—
mt	3.1	2.6	2.6	4.6	3.3	4.4	5.0	5.1	2.6	2.1	4.6	4.1
ne	—	—	—	—	—	—	—	—	—	—	—	0.8
ol	—	—	—	6.9	—	—	—	3.7	3.8	—	—	14.4
il	1.9	1.8	1.4	2.2	2.0	2.5	2.9	3.0	1.1	1.7	2.6	2.6
ap	0.7	0.5	0.4	0.5	0.5	0.7	0.6	0.7	0.3	0.4	0.7	1.0
cc	tr	tr	0.2	0.2	0.3	—	0.7	0.6	—	0.1	0.2	0.2
Total	98.6	98.5	98.8	98.5	98.8	99.1	97.9	98.5	98.9	98.1	98.5	98.2

	Granodiorite of Red Lake			Mount Givens Granodiorite							
	Chinese Peak			Mono Hot Springs		Camp 61					
	Granodiorite	Inclusion		Granodiorite	Inclusion	Granodiorite			Aggregate	Inclusion	
	RL1-H	RL2-I	RL3-I	MHa1-H	MHa2-I	MHb1-H	MHb2-H	MHb3-H	MHb4-A	MHb5-A	MHb6-I	MHb7-I
Q	29.0	—	0.2	22.2	—	23.6	24.6	24.5	8.6	16.0	3.1	1.5
or	17.1	15.5	12.5	16.5	9.5	21.3	21.0	18.2	23.2	19.6	11.8	8.4
ab	30.3	35.6	36.3	27.5	33.1	27.6	26.9	28.4	19.0	27.6	33.0	37.8
an	14.7	23.0	25.3	20.6	25.7	16.8	16.7	17.0	14.6	17.6	23.0	23.8
C	0.8	—	—	—	—	—	—	—	—	—	—	—
di	—	3.5	4.7	0.9	9.8	0.6	0.7	1.3	5.6	3.3	6.8	6.7
hy	3.7	8.2	11.6	6.5	12.5	5.4	5.0	5.0	16.2	8.5	15.4	12.2
mt	1.7	3.9	3.7	2.3	4.5	2.2	2.5	2.5	6.9	3.8	3.0	5.4
ne	—	—	—	—	—	—	—	—	—	—	—	—
ol	—	5.3	—	—	1.0	—	—	—	—	—	—	—
il	0.8	2.1	2.5	1.0	1.8	1.1	1.0	1.2	2.6	1.5	1.9	1.5
ap	0.3	0.7	1.0	0.3	0.5	0.4	0.4	0.4	0.8	0.5	0.6	0.6
cc	0.1	0.1	—	0.1	0.1	0.1	—	—	—	0.1	0.2	—
Total	98.5	97.9	97.8	97.9	98.5	99.1	98.8	98.5	97.5	98.5	98.8	97.9

	Mount Givens Granodiorite—Continued											
	Camp 61—Continued									Courtright Reservoir		
	Inclusion									Granodiorite	Aggregate	
	MHb8-I	MHb9a-Im	MHb9b-Ic	MHb10-I	MHb11-I	MHb12-I	MHb13-I	MHb14-I	MHb15-I	CR1-H	CR2a-A	CR3a-A
Q	—	—	2.3	—	7.5	5.8	—	—	11.7	15.1	15.1	12.2
or	23.2	15.0	9.9	12.9	13.2	8.0	13.3	13.4	8.8	35.2	23.8	20.2
ab	30.5	32.4	34.4	33.4	32.1	35.5	33.7	35.2	34.6	28.7	25.4	21.0
an	18.8	23.6	23.7	24.1	23.1	25.2	22.2	26.6	24.5	12.6	17.0	16.4
C	—	—	—	—	—	—	—	—	—	0.4	—	0.4
di	5.3	6.5	8.2	8.3	1.7	6.4	7.1	2.2	3.0	—	0.6	—
hy	—	—	13.8	0.2	14.1	10.2	9.6	—	10.7	4.7	10.3	18.1
mt	3.6	4.2	3.5	4.2	3.7	4.7	5.5	5.0	3.0	1.2	3.0	4.1
ne	0.1	0.3	—	—	—	—	—	2.7	—	—	—	—
ol	13.2	13.0	—	12.0	—	—	4.3	10.1	—	—	—	—
il	1.9	2.3	2.1	2.4	1.8	1.9	2.0	2.4	1.6	0.9	2.2	3.9
sp	1.0	0.9	0.7	0.9	0.6	0.6	0.6	1.1	0.5	0.2	0.8	1.3
cc	0.1	—	—	—	—	—	—	—	—	—	—	—
Total	97.7	98.2	98.6	98.4	97.8	98.3	98.3	98.7	98.4	99.0	98.2	97.6

Table 4. CIPW norms of mafic inclusions, aggregates, dikes, and their host granitoids, associated gabbroid, and aplite—Continued

	Mount Givens Granodiorite—Continued								
	Courtright Reservoir—Continued								
	Aggregate		Inclusion						Hornfels
	CR4-A	CR5-A	CR2b-I	CR3b-I	CR6-I	CR7-I	CR8-I	CR9-I	CR12-I
Q	13.4	2.8	3.6	10.7	1.2	—	—	0.8	14.9
or	21.6	29.8	12.8	12.1	12.4	12.7	10.1	15.8	0.8
ab	21.6	19.0	35.6	23.8	36.3	38.7	37.0	35.9	8.0
an	16.0	13.5	24.9	18.4	23.9	24.6	23.7	22.2	40.4
C	—	—	—	—	—	—	—	—	—
di	1.3	3.1	0.6	8.3	3.3	5.3	10.0	0.4	30.9
hy	14.7	16.5	13.9	17.7	13.0	4.2	—	15.9	—
mt	4.3	7.6	4.0	4.3	5.3	5.1	4.7	3.9	1.5
ne	—	—	—	—	—	—	0.2	—	—
ol	—	—	—	—	—	5.9	9.8	—	—
il	3.4	4.5	2.1	1.8	2.1	2.3	2.3	2.1	2.0
ap	1.3	1.6	0.9	0.3	0.8	1.0	0.8	0.7	0.8
cc	0.2	—	0.2	0.2	0.1	0.1	—	0.2	—
Total	97.8	98.4	98.6	97.6	98.4	99.9	98.6	97.9	99.3

Table 5. Average chemical analyses, in weight percent, of biotite in mafic inclusions, dikes, aggregates, and their host granitoids

[Chemical analyses by B. Barbarin using an electron microprobe; n, number of analyzed spots included in average; at., atomic; FeO*, total iron as FeO; m, margin; c, core]

Unit	Bass Lake Tonalite								Dinkey Creek Granodiorite			
Locality	North Fork (NF)					Meadow Lakes (ML)			Shaver Lake quarry (SQ)			
Sample No.	1	2	3	4a	4b	1		2	1			4a
Description	H	I	I	D	D	H	I	I	H	A	I	H
n	6	6	7	5	5	4	4	6	5	6	5	4
SiO_2	35.1	34.8	36.7	36.4	37.2	36.9	37.0	37.2	36.0	36.0	36.1	36.0
Al_2O_3	16.1	15.8	15.4	15.6	15.8	15.7	15.4	15.8	15.3	14.8	14.7	15.6
FeO*	20.8	21.0	21.1	20.0	19.3	20.9	20.9	20.9	22.3	22.4	22.8	21.7
MgO	9.78	9.70	9.94	10.24	11.25	9.87	9.51	10.18	9.24	9.87	9.54	9.65
CaO	0.04	0.04	0.04	0.02	0.03	0.02	0.03	0.03	0.01	0.02	0.01	0.02
Na_2O	0.08	0.09	0.09	0.14	0.09	0.09	0.10	0.09	0.07	0.07	0.07	0.07
K_2O	9.47	9.34	9.57	9.52	9.42	9.50	9.45	9.46	9.43	9.45	9.48	9.66
TiO_2	3.17	3.38	3.21	3.93	3.14	3.07	3.53	2.91	3.45	3.25	3.47	3.13
MnO	0.30	0.30	0.28	0.12	0.28	0.29	0.29	0.22	0.35	0.34	0.34	0.31
Total	94.8	94.5	96.3	96.0	96.5	96.3	96.2	96.8	96.2	96.2	96.5	96.1
at.Fe/Fe+Mg	0.54	0.55	0.54	0.52	0.49	0.54	0.55	0.54	0.58	0.56	0.57	0.56

Unit	Dinkey Creek Granodiorite–Continued											
Locality	Shaver Lake quarry (SQ)–Continued											
Sample No.	4a–Continued		4b		5		6		9		10	
Description	A	I	I(dark)	I(light)	H	I	H	I	H	I	H	I
n	9	3	4	4	3	7	4	6	3	7	4	7
SiO_2	36.2	36.5	35.7	35.7	36.5	36.6	36.3	36.3	36.3	36.6	36.2	36.5
Al_2O_3	15.6	15.7	15.5	15.8	15.7	15.3	15.1	15.2	15.6	15.8	15.7	16.0
FeO*	21.6	21.3	20.8	20.8	21.5	21.4	21.4	21.4	21.7	21.5	21.5	21.4
MgO	9.75	9.93	9.94	10.20	9.40	9.79	9.35	9.74	9.68	9.86	9.46	9.90
CaO	0.03	0.02	0.04	0.02	0.03	0.03	0.01	0.04	0.03	0.06	0.03	0.05
Na_2O	0.07	0.08	0.07	0.07	0.08	0.09	0.08	0.07	0.08	0.08	0.07	0.08
K_2O	9.72	9.86	9.56	9.57	9.61	9.50	9.39	9.36	9.33	9.24	9.30	9.25
TiO_2	3.17	3.05	3.36	2.74	3.20	2.83	3.35	3.08	2.95	2.88	3.20	2.50
MnO	0.31	0.33	0.31	0.30	0.33	0.34	0.33	0.33	0.34	0.35	0.34	0.34
Total	96.5	96.8	95.3	95.2	96.4	95.9	95.3	95.5	96.0	96.4	95.8	96.0
at.Fe/Fe+Mg	0.55	0.55	0.54	0.53	0.56	0.55	0.56	0.55	0.56	0.55	0.56	0.55

Unit	Dinkey Creek Granodiorite–Continued											
Locality	Shaver lake quarry (SQ)–Continued				Markwood Creek (MK)			Deer Creek (DR)				
Sample No.	12		13		1	2		1	2	3	4	
Description	H	I	H	I	H	H	I	H	A	I	H	I
n	5	5	4	6	5	7	5	7	9	6	5	7
SiO_2	36.1	36.1	36.2	36.4	36.0	36.4	36.3	36.3	36.9	37.2	36.9	37.3
Al_2O_3	15.7	15.7	15.7	15.6	15.9	16.0	15.7	15.6	15.2	16.0	15.4	15.7
FeO*	21.7	21.6	21.8	21.6	22.2	22.4	22.2	20.4	20.1	19.6	19.6	18.0
MgO	9.50	9.47	9.35	9.54	9.35	9.37	9.5	10.64	10.84	11.27	11.17	12.11
CaO	0.01	0.02	0.01	0.02	0.03	0.01	0.01	0.01	0.03	0.02	0.03	0.05
Na_2O	0.07	0.07	0.08	0.08	0.08	0.08	0.08	0.15	0.16	0.14	0.15	0.14
K_2O	9.75	9.76	9.78	9.78	9.39	9.58	9.55	9.44	9.42	9.47	9.50	9.55
TiO_2	3.23	3.46	3.63	3.18	3.59	3.50	4.16	3.62	3.34	2.76	3.24	2.96
MnO	0.35	0.36	0.33	0.34	0.32	0.29	0.31	0.26	0.22	0.16	0.15	0.09
Total	96.4	96.5	96.9	96.5	96.9	97.6	97.8	96.4	96.2	96.6	96.1	95.9
at.Fe/Fe+Mg	0.56	0.56	0.57	0.56	0.57	0.57	0.57	0.52	0.51	0.49	0.50	0.46

Table 5. Average chemical analyses, in weight percent, of biotite in mafic inclusions, dikes, aggregates, and their host granitoids—Continued

Unit	Dinkey Creek Granodiorite–Continued										Granodiorite of Red lake	
Locality	Deer Creek (DR)–Continued					Wishon Reservoir (WR)					Chinese Peak (CP)	
Sample No.	5			6		1	2		3		1	
Description	H	I	I	H	I	I	A	I	H	I	H	I(c)
n	5	3	4	5	5	4	6	4	4	7	5	5
SiO_2	36.9	37.0	37.3	36.5	36.9	36.2	37.0	36.9	37.0	37.2	37.0	37.1
Al_2O_3	16.2	16.4	16.0	16.1	16.0	15.8	15.9	15.7	15.8	15.8	14.9	14.6
FeO*	19.1	18.3	19.2	18.8	18.4	20.3	21.3	21.1	21.6	21.2	20.7	20.4
MgO	11.44	12.04	11.38	11.97	12.10	11.06	10.50	10.69	10.23	10.45	10.47	10.50
CaO	0.02	0.05	0.02	0.03	0.02	0.03	0.02	0.03	0.03	0.01	0.01	0.01
Na_2O	0.17	0.12	0.12	0.16	0.17	0.09	0.08	0.08	0.08	0.08	0.10	0.10
K_2O	9.42	9.64	9.59	9.54	9.54	9.46	9.32	9.23	9.22	9.30	9.17	9.23
TiO_2	2.88	2.26	3.00	2.49	3.10	2.96	2.48	2.28	2.70	2.56	3.77	3.88
MnO	0.14	0.11	0.13	0.13	0.10	0.37	0.30	0.29	0.28	0.29	0.48	0.42
Total	96.3	95.9	96.7	95.7	96.3	96.3	96.9	96.3	96.9	96.9	96.6	96.2
at.Fe/Fe+Mg	0.48	0.46	0.49	0.47	0.46	0.51	0.53	0.53	0.54	0.53	0.53	0.52

Unit	Granodiorite of Red Lake–Continued		Mount Givens Granodiorite									
Locality	Chinese Peak (CP)–Continued		Mono Hot Springs (MHa)			Camp 61 (MHb)						
Sample No.	1	2	1	2		1	6	7	8	9		
Description	I(m)	I	H	H	I	H	I	I	I	H	I(c)	I(m)
n	6	5	4	4	6	7	6	5	4	3	5	6
SiO_2	37.3	36.4	37.3	36.8	36.8	38.3	35.8	36.2	36.2	35.9	36.3	36.1
Al_2O_3	14.8	14.7	15.8	16.0	15.7	15.6	15.1	15.5	15.5	15.5	15.4	15.4
FeO*	20.0	20.7	19.9	21.1	21.4	18.6	18.6	18.7	17.9	18.7	18.7	18.7
MgO	10.79	10.60	11.2	10.4	10.4	12.4	11.6	11.7	12.6	11.6	11.7	11.7
CaO	0.03	0.03	0.02	0.02	0.02	0.03	0.01	0.01	0.05	0.02	0.03	0.04
Na_2O	0.11	0.10	0.08	0.08	0.09	0.09	0.08	0.09	0.07	0.08	0.08	0.08
K_2O	9.25	9.66	9.26	9.25	9.17	9.49	9.47	9.47	9.66	9.45	9.55	9.45
TiO_2	3.62	3.68	2.72	2.42	2.51	2.37	3.23	2.43	1.93	2.87	3.06	2.80
MnO	0.43	0.40	0.44	0.29	0.30	0.47	0.39	0.46	0.35	0.41	0.35	0.39
Total	96.3	96.3	96.7	96.4	96.4	97.4	94.3	94.6	94.3	94.5	95.2	94.7
at.Fe/Fe+Mg	0.51	0.52	0.50	0.54	0.54	0.46	0.47	0.47	0.45	0.48	0.47	0.47

Unit	Mount Givens Granodiorite–Continued						
Locality	Camp 61 (MHb)–Continued						
Sample No.	10			11		12	
Description	H	I(c)	I(m)	H	I	H	I
n	4	6	4	6	5	4	7
SiO_2	36.2	36.3	36.3	36.7	36.8	36.8	36.8
Al_2O_3	15.3	15.6	15.7	15.5	15.4	15.4	15.3
FeO*	18.7	18.7	18.3	18.8	18.8	18.7	18.8
MgO	11.8	11.8	11.9	11.7	11.5	11.7	11.8
CaO	0.02	0.04	0.06	0.02	0.02	0.01	0.03
Na_2O	0.07	0.07	0.07	0.08	0.08	0.09	0.08
K_2O	9.51	9.48	9.37	9.58	9.59	9.66	9.54
TiO_2	2.85	2.72	2.35	2.62	3.11	2.61	2.67
MnO	0.37	0.38	0.41	0.42	0.39	0.43	0.39
Total	94.8	95.1	94.5	95.4	95.7	95.4	95.4
at.Fe/Fe+Mg	0.47	0.47	0.46	0.47	0.48	0.47	0.47

Table 6. Average chemical analyses, in weight percent, of hornblende in mafic inclusions, dikes, aggregates, and their host granitoids and associated gabbroid

[Chemical analyses by B. Barbarin using an electron microprobe; n, number of analyzed spots included in average; at., atomic; FeO*, total iron as FeO; m, margin; c, core]

Unit	Bass Lake Tonalite								Dinkey Creek Granodiorite			
Locality	North Fork (NF)					Meadow Lakes (ML)			Shaver Lake quarry (SQ)			
Sample No.	1	2	3	4a	5	1		2	1			4a
Description	H	I	I	D	G	H	I	I	H	A	I	H
n	4	6	6	5	5	5	3	5	5	4	6	4
SiO_2	44.2	43.5	46.3	46.5	46.7	45.4	45.6	45.6	44.6	44.5	44.3	44.3
Al_2O_3	8.77	8.81	8.35	8.85	8.48	8.69	8.75	8.61	8.33	8.31	8.39	8.25
FeO*	18.1	18.2	18.6	16.1	16.6	18.5	18.9	18.5	19.7	19.8	20.3	19.2
MgO	10.07	9.87	10.08	11.31	11.50	9.99	9.68	10.01	9.70	9.67	9.34	9.82
CaO	12.5	12.7	12.4	12.3	12.1	12.2	12.4	12.4	11.9	11.8	12.0	12.9
Na_2O	1.07	1.01	1.00	1.09	0.99	1.09	0.97	1 01	1.13	1.30	1.18	1.04
K_2O	0.75	0.84	0.82	0.44	0.83	0.88	0.93	0.89	0.92	0.91	0.94	0.86
TiO_2	1.23	0.96	0.91	1.27	1.11	1.43	1.24	1.45	1.39	1.54	1.33	1.28
MnO	0.51	0.51	0.50	0.29	0.49	0.46	0.44	0.43	0.53	0.52	0.54	0.50
Total	97.2	96.4	99.0	98.2	98.8	98.6	98.9	98.9	98.2	98.4	98.3	98.2
at.Fe/Fe+Mg	0.50	0.51	0.51	0.45	0.45	0.51	0.52	0.51	0.53	0.54	0.55	0.52

Unit	Dinkey Creek Granodiorite—Continued											
Locality	Shaver Lake quarry (SQ)—Continued											
Sample No.	4—Continued				5		6		9		10	
Description	A	I	I (dark)	I (light)	H	I	H	I	H	I	H	I
n	7	3	6	4	4	7	4	6	3	7	4	7
SiO_2	44.8	45.1	44.1	44.7	45.3	44.6	45.0	44.8	44.5	45.1	44.2	44.4
Al_2O_3	8.31	8.52	8.13	8.16	8.08	8.66	8.21	8.32	8.48	8.49	8.67	8.91
FeO*	18.9	19.2	18.4	18.1	19.0	19.6	19.0	19.4	19.1	19.3	19.5	19.7
MgO	10.00	9.89	10.23	10.40	9.91	9.33	9.73	9.49	9.81	9.71	9.22	9.30
CaO	12.6	12.6	13.2	13.0	11.9	12.0	12.1	12.2	11.8	12.4	12.1	12.4
Na_2O	1.12	1.02	0.97	0.98	1.10	1.11	1.09	1.02	1.17	1.01	1.07	0.99
K_2O	0.88	0.90	0.83	0.85	0.88	0.99	0.89	0.91	0.92	0.89	0.95	0.92
TiO_2	1.44	1.19	1.19	1.21	1.30	1.31	1.38	1.24	1.43	1.20	1.32	1.19
MnO	0.50	0.49	0.49	0.47	0.53	0.53	0.50	0.49	0.48	0.54	0.53	0.51
Total	98.6	98.9	97.5	97.9	98.0	98.1	97.9	97.9	97.7	98.6	97.6	98.30
at.Fe/Fe+Mg	0.51	0.52	0.50	0.49	0.52	0.54	0.52	0.54	0.52	0.53	0.54	0.54

Unit	Dinkey Creek Granodiorite—Continued											
Locality	Shaver lake quarry (SQ)—Continued				Markwood Creek (MK)			Deer Creek (DR)				
Sample No.	12		13		1	2		1	2	3	4	
Description	H	I	H	I	H	H	I	H	A	I	H	I
n	5	6	3	7	5	5	7	7	9	6	5	7
SiO_2	44.2	44.3	44.7	44.5	45.1	44.6	44.2	45.4	46.0	47.1	45.9	45.5
Al_2O_3	8.68	8.70	8.35	8.68	8.51	8.87	9.32	8.52	8.14	8.27	8.70	9.47
FeO	19.7	19.6	19.6	19.6	19.8	20.1	20.0	17.8	15.6	15.0	16.2	16.1
MgO	9.34	9.35	9.53	9.43	9.72	9.52	9.37	10.89	10.95	13.01	12.12	11.81
CaO	11.8	11.9	11.7	11.8	12.0	12.0	12.1	11.9	12.4	11.9	11.7	11.8
Na_2O	1.13	1.06	1.15	1.06	1.06	1.12	1.10	1.09	1.03	0.99	1.06	1.05
K_2O	0.97	0.95	0.95	0.96	0.92	1.00	1.05	0.83	0.82	0.63	0.63	0.59
TiO_2	1.36	1.20	1.32	1.27	1.39	1.37	1.38	1.48	1.49	1.65	1.64	1.69
MnO	0.53	0.53	0.52	0.52	0.49	0.49	0.48	0.46	0.43	0.31	0.32	0.28
Total	97.7	97.6	97.8	97.8	99.0	99.1	99.0	98.4	96.9	98.9	98.3	98.3
at.Fe/Fe+Mg	0.54	0.54	0.54	0.54	0.53	0.54	0.55	0.48	0.47	0.39	0.43	0.43

Table 6. Average chemical analyses, in weight percent, of hornblende in mafic inclusions, dikes, aggregates, and their host granitoids and associated gabbroid—Continued

Unit	Dinkey Creek Granodiorite—Continued										Granodiorite of Red lake	
Locality	Deer Creek (DR)—Continued					Wishon Reservoir (WR)					Chinese Peak (CP)	
Sample No.	5			6		1	2		3		1	
Description	H	I	I	H	I	I	A	I	H	I	H	I(c)
n	5	3	4	5	5	4	6	4	4	7	4	5
SiO_2	46.6	46.2	46.8	45.4	46.3	46.0	46.4	46.0	46.3	46.5	46.4	46.3
Al_2O_3	8.94	9.25	8.75	9.09	8.64	7.87	8.32	8.75	8.57	8.44	8.26	8.07
FeO	16.1	16.1	16.1	16.2	16.1	17.7	18.4	18.9	18.4	18.5	16.8	17.3
MgO	12.13	11.94	12.15	12.02	12.30	11.21	10.54	10.11	10.42	10.38	11.54	11.07
CaO	12.5	11.5	11.4	11.2	11.2	12.3	12.4	12.5	12.3	12.5	12.1	12.2
Na_2O	0.98	0.96	0.95	1.05	0.93	0.99	0.97	0.92	1.01	0.96	1.36	1.27
K_2O	0.63	0.60	0.54	0.75	0.56	0.75	0.83	0.85	0.81	0.80	0.81	0.80
TiO_2	1.56	1.64	1.38	1.80	1.41	1.20	1.42	1.08	1.28	1.19	1.30	1.29
MnO	0.32	0.29	0.37	0.30	0.33	0.55	0.47	0.46	0.46	0.47	0.58	0.65
Total	99.8	98.5	98.4	97.8	97.8	98.6	99.8	99.6	99.6	99.7	99.2	99.0
at.Fe/Fe+Mg	0.43	0.43	0.43	0.43	0.42	0.47	0.50	0.51	0.50	0.50	0.45	0.47

Unit	Granodiorite of Red Lake—Continued		Mount Givens Granodiorite									
Locality	Chinese Peak (CP)—Continued		Mono Hot Springs (MHa)			Camp 61 (MHb)						
Sample No.	1	2	1	2		1	6	7	8	9		
Description	I(m)	I	H	I	I	H	I	I	I	H	I(c)	I(m)
n	5	5	4	4	6	5	6	5	5	3	6	9
SiO_2	47.6	46.5	46.8	45.8	45.7	48.4	45.3	45.9	47.5	45.4	45.1	45.2
Al_2O_3	7.31	7.25	8.11	8.80	8.64	7.42	7.28	7.19	6.07	7.84	7.81	7.98
FeO*	16.5	16.5	17.1	18.9	18.8	16.1	15.8	15.9	14.8	16.0	16.2	16.4
MgO	11.96	11.93	11.5	10.0	10.0	12.4	12.1	12.1	13.5	12.0	11.9	11.6
CaO	12.1	11.7	12.3	12.4	12.4	12.6	13.3	13.1	12.7	13.0	13.2	13.2
Na_2O	1.20	1.29	1.04	1.01	0.97	1.05	0.96	0.99	0.78	1.10	1.06	1.01
K_2O	0.67	0.70	0.78	0.85	0.82	0.71	0.67	0.69	0.51	0.81	0.79	0.78
TiO_2	1.23	1.26	1.28	1.22	1.27	1.10	1.14	1.22	0.65	1.32	1.23	1.18
MnO	0.61	0.58	0.59	0.47	0.47	0.64	0.60	0.63	0.54	0.58	0.56	0.58
Total	99.2	97.7	99.5	99.5	99.1	100.4	97.2	97.7	97.1	98.1	97.9	97.9
at.Fe/Fe+Mg	0.44	0.44	0.46	0.52	0.51	0.42	0.42	0.43	0.38	0.43	0.43	0.44

Unit	Mount Givens Granodiorite—Continued						
Locality	Camp 61 (MHb)—Continued						
Sample No.	10			11		12	
Description	H	I(c)	I(m)	H	I	H	I
n	4	4	6	5	6	5	7
SiO_2	45.1	45.5	45.9	47.0	46.7	46.6	46.5
Al_2O_3	7.90	7.64	7.52	7.26	7.95	7.49	7.42
FeO*	15.7	16.5	16.1	15.8	15.8	15.9	16.2
MgO	12.2	11.6	12.0	12.4	12.2	12.3	12.1
CaO	13.0	13.3	13.1	12.2	12.2	12.1	12.1
Na_2O	1.15	0.97	0.95	0.94	0.94	1.04	0.98
K_2O	0.79	0.71	0.68	0.69	0.71	0.74	0.74
TiO_2	1.29	1.02	1.12	1.00	1.14	1.07	1.03
MnO	0.53	0.56	0.56	0.60	0.61	0.57	0.57
Total	97.7	97.8	97.9	97.9	98.3	97.8	97.6
at.Fe/Fe+Mg	0.42	0.44	0.43	0.42	0.42	0.42	0.43

Table 7. Rb-Sr isotopic data on mafic inclusions, dikes, aggregates, and their host granitoids

[Rb and Sr concentrations by energy dispersive X-ray fluorescence. Analyses by A.C. Robinson and M. Wilcox. c, coarse; f, fine]

Sample	Rb(ppm)	Sr(ppm)	$^{87}Rb/^{86}Sr$	$^{87}Sr/^{86}Sr$
Bass Lake Tonalite				
HH1-H	103	360	0.83	0.70723±6
HH2-A	65.3	488	0.39	0.70650±6
HH3-A	68.8	455	0.44	0.70677±6
HH4-I	86.1	486	0.51	0.70676±5
HH5a-D	74.1	665	0.32	0.70653±3
HH5b-D	63.8	654	0.28	0.70640±3
HH5c-D	61.3	619	0.29	0.70608±4
HH5d-D	54.9	595	0.27	0.70643±3
HH6a-D	58.1	580	0.29	0.70566±4
HH6b-D	61.8	569	0.31	0.70565±5
HH6c-D	54.4	637	0.25	0.70652±5
HH6d-D	68.8	525	0.38	0.70612±3
HH6e-AP	90.1	385	0.68	0.70684±6
NF1-H	66.5	276	0.70	0.70562±4
NF2-I	61.7	245	0.73	0.70573±4
NF3-I	87.8	253	1.00	0.70616±4
NF4a-D	34.3	769	0.13	0.70633±3
NF4b-D	40.6	620	0.19	0.70565±4
NF5-G	43.9	450	0.28	0.70541±4
Dinkey Creek Granodiorite				
SQ1-H	132	310	1.23	0.70930±4
SQ2-H	125	330	1.10	0.70914±5
SQ3-I	103	336	0.89	0.70886±5
SQ3a-A	161	207	2.25	0.71089±5
SQ3b-Ac	196	180	3.15	0.71215±5
SQ3b-Af	156	276	1.64	0.70980±6
SQ4-I	101	328	0.89	0.70881±5
SQ5-I	142	292	1.41	0.70945±4
SQ6-I	110	321	0.99	0.70893±5
SQ7-I	96.3	344	0.81	0.70874±4
SQ8-I	161	321	1.45	0.70966±4
SQ9-I	110	281	1.13	0.70916±5
SQ11-I	169	330	1.48	0.70969±3
DR1a-H	62.6	471	0.38	0.70787±5
Dinkey Creek Granodiorite—Continued				
DR2-A	76.6	382	0.58	0.70797±3
DR3-I	50.6	471	0.31	0.70744±4
DR4a-I	15.4	578	0.08	0.70745±4
DR5-I	13.1	575	0.07	0.70739±3
WR1-H	96.1	402	0.69	0.70834±3
WR2-A	124	257	1.40	0.70920±4
WR3-I	124	380	0.94	0.70852±3
Mount Givens Granodiorite				
MHb1-H	130	414	0.91	0.70765±4
MHb2-H	129	425	0.88	0.70762±3
MHb3-H	126	434	0.84	0.70757±8
MHb4-A	207	262	2.29	0.70942±6
MHb5-A	142	387	1.06	0.70781±3
MHb6-I	141	356	1.15	0.70783±5
MHb7-I	77.0	403	0.55	0.70717±4
MHb8-I	291	360	2.34	0.70925±3
MHb9a-Im	167	362	1.33	0.70814±5
MHb9b-Ic	103	364	0.82	0.70731±5
MHb10-I	172	375	1.33	0.70816±5
MHb13-I	210	329	2.43	0.70875±8
MHb14-I	193	410	1.36	0.70819±6
MHb15-I	98.1	403	0.70	0.70737±3
CR1-H	184	484	1.10	0.70807±5
CR2a-A	120	533	0.65	0.70734±3
CR2b-I	104	553	0.54	0.70721±3
CR3a-A	163	390	1.21	0.70805±5
CR3b-I	84.8	352	0.70	0.70733±3
CR4-A	141	395	1.03	0.70780±4
CR5-A	286	330	2.51	0.70966±4
CR6-I	115	522	0.64	0.70732±4
CR7-I	115	516	0.64	0.70735±5
CR8-I	101	510	0.57	0.70727±5
CR9-I	175	512	0.99	0.70774±6
CR10-I	0.2	275	0.003	0.70987±1

Table 8. Sm-Nd isotopic data on mafic inclusions, aggregates, and associated rocks

[Sm and Nd concentrations from table 3. Data for SLb96X3, host granitoid, from Domenick and others (1983)]

Sample No.	Sm (ppm)	Nd (ppm)	$^{147}Sm/^{144}Nd$	$^{143}Nd/^{144}Nd$
Bass Lake Tonalite				
HH1-H	1.3	13.9	0.057	0.51231±3
HH4-I	5.9	34.6	0.103	0.51237±2
HH6a-D	1.8	25.3	0.043	0.51246±2
Dinkey Creek Granodiorite				
Slb96X3	5.6	32.2	0.105	0.51218±1
SQ2-H	3.9	21.9	0.107	0.51220±2
SQ5-I	8.1	39.0	0.126	0.51225±2
SQ9-I	5.6	25.0	0.135	0.51233±2
DR2-A	6.7	26.9	0.151	0.51229±3
Mount Givens Granodiorite				
MHb3-H	3.8	34.1	0.067	0.51237±2
MHb4-A	10.1	56.6	0.108	0.51246±2
MHb13-I	6.1	28.4	0.130	0.51248±2

SELECTED SERIES OF U.S. GEOLOGICAL SURVEY PUBLICATIONS

Periodicals

Earthquakes & Volcanoes (issued bimonthly).
Preliminary Determination of Epicenters (issued monthly).

Technical Books and Reports

Professional Papers are mainly comprehensive scientific reports of wide and lasting interest and importance to professional scientists and engineers. Included are reports on the results of resource studies and of topographic, hydrologic, and geologic investigations. They also include collections of related papers addressing different aspects of a single scientific topic.

Bulletins contain significant data and interpretations that are of lasting scientific interest but are generally more limited in scope or geographic coverage than Professional Papers. They include the results of resource studies and of geologic and topographic investigations; as well as collections of short papers related to a specific topic.

Water-Supply Papers are comprehensive reports that present significant interpretive results of hydrologic investigations of wide interest to professional geologists, hydrologists, and engineers. The series covers investigations in all phases of hydrology, including hydrogeology, availability of water, quality of water, and use of water.

Circulars present administrative information or important scientific information of wide popular interest in a format designed for distribution at no cost to the public. Information is usually of short-term interest.

Water-Resources Investigations Reports are papers of an interpretive nature made available to the public outside the formal USGS publications series. Copies are reproduced on request unlike formal USGS publications, and they are also available for public inspection at depositories indicated in USGS catalogs.

Open-File Reports include unpublished manuscript reports, maps, and other material that are made available for public consultation at depositories. They are a nonpermanent form of publication that may be cited in other publications as sources of information.

Maps

Geologic Quadrangle Maps are multicolor geologic maps on topographic bases in 7 1/2- or 15-minute quadrangle formats (scales mainly 1:24,000 or 1:62,500) showing bedrock, surficial, or engineering geology. Maps generally include brief texts; some maps include structure and columnar sections only.

Geophysical Investigations Maps are on topographic or planimetric bases at various scales; they show results of surveys using geophysical techniques, such as gravity, magnetic, seismic, or radioactivity, which reflect subsurface structures that are of economic or geologic significance. Many maps include correlations with the geology.

Miscellaneous Investigations Series Maps are on planimetric or topographic bases of regular and irregular areas at various scales; they present a wide variety of format and subject matter. The series also includes 7 1/2-minute quadrangle photogeologic maps on planimetric bases which show geology as interpreted from aerial photographs. Series also includes maps of Mars and the Moon.

Coal Investigations Maps are geologic maps on topographic or planimetric bases at various scales showing bedrock or surficial geology, stratigraphy, and structural relations in certain coal-resource areas.

Oil and Gas Investigations Charts show stratigraphic information for certain oil and gas fields and other areas having petroleum potential.

Miscellaneous Field Studies Maps are multicolor or black-and-white maps on topographic or planimetric bases on quadrangle or irregular areas at various scales. Pre-1971 maps show bedrock geology in relation to specific mining or mineral-deposit problems; post-1971 maps are primarily black-and-white maps on various subjects such as environmental studies or wilderness mineral investigations.

Hydrologic Investigations Atlases are multicolored or black-and-white maps on topographic or planimetric bases presenting a wide range of geohydrologic data of both regular and irregular areas; principal scale is 1:24,000 and regional studies are at 1:250,000 scale or smaller.

Catalogs

Permanent catalogs, as well as some others, giving comprehensive listings of U.S. Geological Survey publications are available under the conditions indicated below from the U.S. Geological Survey, Books and Open-File Reports Section, Federal Center, Box 25425, Denver, CO 80225. (See latest Price and Availability List.)

"Publications of the Geological Survey, 1879-1961" may be purchased by mail and over the counter in paperback book form and as a set of microfiche.

"Publications of the Geological Survey, 1962-1970" may be purchased by mail and over the counter in paperback book form and as a set of microfiche.

"Publications of the U.S. Geological Survey, 1971-1981" may be purchased by mail and over the counter in paperback book form (two volumes, publications listing and index) and as a set of microfiche.

Supplements for 1982, 1983, 1984, 1985, 1986, and for subsequent years since the last permanent catalog may be purchased by mail and over the counter in paperback book form.

State catalogs, "List of U.S. Geological Survey Geologic and Water-Supply Reports and Maps For (State)," may be purchased by mail and over the counter in paperback booklet form only.

"Price and Availability List of U.S. Geological Survey Publications," issued annually, is available free of charge in paperback booklet form only.

Selected copies of a monthly catalog "New Publications of the U.S. Geological Survey" available free of charge by mail or may be obtained over the counter in paperback booklet form only. Those wishing a free subscription to the monthly catalog "New Publications of the U.S. Geological Survey" should write to the U.S. Geological Survey, 582 National Center, Reston, VA 22092.

Note.--Prices of Government publications listed in older catalogs, announcements, and publications may be incorrect. Therefore, the prices charged may differ from the prices in catalogs, announcements, and publications.